


IL-36 γ drives skin toxicity induced by EGFR/MEK inhibition and commensal *Cutibacterium acnes*

Takashi K. Satoh, ... , Emmanuel Contassot, Lars E. French

J Clin Invest. 2019. <https://doi.org/10.1172/JCI128678>.

Research In-Press Preview Dermatology Inflammation

Graphical abstract



Find the latest version:

<https://jci.me/128678/pdf>



IL-36 γ drives skin toxicity induced by EGFR/MEK inhibition and commensal *Cutibacterium acnes*

Authors: Takashi K. Satoh^{1*}, Mark Mellett¹, Barbara Meier-Schiesser¹, Gabriele Fenini¹, Atsushi Otsuka², Hans-Dietmar Beer^{1,4}, Tamara Rordorf³, Julia-Tatjana Maul¹, Jürg Hafner^{1,4}, Alexander A. Navarini^{1,4,5}, Emmanuel Contassot^{1,4†}, Lars E. French^{1,4,6*†}

Affiliations:

¹Department of Dermatology, University of Zürich, Zürich, Switzerland.

²Department of Dermatology, Kyoto University, Kyoto, Japan.

³Clinic for Oncology, University Hospital Zürich, Zürich, Switzerland.

⁴Medical Faculty, University of Zürich, Zürich, Switzerland.

⁵Department of Dermatology, University Hospital of Basel, Basel, Switzerland

⁶Department of Dermatology and Allergology, Ludwig Maximilian University of Munich, Munich, Germany.

* co-corresponding authors

† co-senior authors

co-corresponding authors:

Takashi K. Satoh: Department of Dermatology, University of Zurich, Gloriastrasse 31, Switzerland, 8091, +41 (0) 44 255 4115, Takashi.Satoh@usz.ch

Lars E. French: Department of Dermatology and Allergology, Ludwig-Maximilians-University of Munich, Frauenlobstrasse 9–11, Germany, 80337, +49 (89) 44005 6005, Lars.French@med.uni-muenchen.de

Abstract: Epidermal growth factor receptor (EGFR) and MEK inhibitors (EGFR/MEKi) are beneficial for the treatment of solid cancers but are frequently associated with severe therapy-limiting acneiform skin toxicities. The underlying molecular mechanisms are poorly understood. Using gene expression profiling we identified IL-36 γ and IL-8 as candidate drivers of EGFR/MEKi skin toxicity. We provide molecular and translational evidence that EGFR/MEKi in concert with the skin commensal bacterium *Cutibacterium acnes* act synergistically to induce IL-36 γ in keratinocytes and subsequently IL-8, leading to cutaneous neutrophilia. IL-36 γ expression was the combined result of *C. acnes*-induced NF- κ B activation and EGFR/MEKi-mediated expression of the transcription factor Krüppel-like factor 4 (KLF4), due to the presence of both NF- κ B- and KLF4-binding sites in the human IL-36 γ gene promoter. EGFR/MEKi increased KLF4 expression by blockade of the EGFR-MEK-ERK pathway. These results provide an insight into understanding the pathological mechanism of the acneiform skin toxicities induced by EGFR/MEKi and identify IL-36 γ and the transcription factor KLF4 as potential therapeutic targets.

Introduction

Agents targeting the epidermal growth factor receptor (EGFR)-mediated signaling pathway are increasingly used for the treatment of advanced lung, pancreatic, colorectal and head and neck cancers, which benefit from exacerbated EGFR activity for their growth and survival (1, 2). Small molecule inhibitors of the Mitogen-activated protein kinases (MAPKs) signaling pathways including extracellular signal-regulated kinase (ERK) and MAPK/ERK kinase (MEK) 1 and 2 have also shown their efficacy in the treatment of various cancers, especially melanoma (3, 4).

One significant clinical limitation to the prolonged use of EGFR and MEK inhibitors (EGFR/MEKi) is the occurrence of skin toxicities in 50-80% of patients, including an acneiform eruption that usually develops within the first few weeks of therapy (4-6). The first monoclonal antibody and small molecule inhibitor of EGFR were approved by the FDA for the treatment of cancer twelve and thirteen years ago respectively, and since then skin toxicities induced by these drugs remain unsolved problems. Even though topical or oral agents alone or in combination are used to treat skin toxicities and show some efficacy, EGFR/MEKi-induced skin acneiform eruptions can still seriously affect patients' quality of life, leading physicians to reduce the dose administered or discontinue therapy in severe skin toxicity cases (7-9). Importantly, the development and severity of the acneiform eruption have been shown to correlate with favorable anti-tumor responses (10-12).

The exact molecular pathogenesis underlying the frequent and rapid development of skin toxicity to EGFR/MEKi is not understood to date. Animal studies using mice selectively lacking EGFR in the skin revealed that EGFR signaling is critical for normal skin barrier function and antimicrobial defense (13, 14). However, the phenotype of mice selectively lacking EGFR in the skin resembles atopic dermatitis and is distinct from the acneiform skin toxicity seen in patients treated with EGFR/MEKi (15, 16). In humans, histopathology of acneiform eruption lesions is characterized by folliculitis with massive infiltration of neutrophils histologically resembling acne vulgaris (5). Another feature of acneiform toxicities caused by EGFR/MEKi and shared with acne vulgaris is the topographical predominance of inflammation in skin areas rich in pilosebaceous units, also corresponding to sebum-rich regions of the skin, such as the central face, upper chest, and back (17-21). These sebum-rich regions are highly colonized by *Cutibacterium acnes* (*C. acnes*, formerly known as *Propionibacterium acnes*) a lipophilic commensal representing the most abundant microorganism on the skin of healthy adults (19-22). While *C. acnes* is thought to play an

important role in common acne, its involvement in EGFR/MEKi acneiform toxicities has never been investigated.

A better understanding of the molecular pathogenesis of acneiform eruption by EGFR/MEKi is still needed so as to guide the development of effective therapies to prevent or suppress the skin toxicity, while preserving their anti-tumor effects. Here, we investigate the molecular mechanisms of acneiform eruption associated with EGFR/MEKi.

Results

Skin gene expression profiling in EGFRi-induced acneiform skin toxicity

Employing an unbiased approach, we performed gene expression profiling of lesional skin biopsy samples from patients suffering from acneiform eruption by EGFRi (Figure 1A and Supplemental Table 1). We found elevated IL-8 and IL-36 γ in the patients' skin, whereas important inflammatory cytokines such as TNF- α , IL-6 and IL-17A were not significantly upregulated when compared to skin from healthy donors (Figure 1A). This observation was further confirmed by quantitative PCR with more lesional skin samples (Figure 1B and Supplemental Figure 1A). As previously reported, the expression of antimicrobial peptides such as RNase7 was also found to be decreased in patients' skin (14) (Supplemental Figure 1A). IL-36 γ is a proinflammatory cytokine of the IL-1 family, predominantly expressed by keratinocytes and is able to signal in an auto- or paracrine manner through the IL-36 receptor (also known as IL1RL2) and activates NF- κ B signaling pathway in target cells. It has recently been shown that IL-36 plays a role in the cutaneous neutrophilic pustular autoinflammatory disease called DITRA (Deficiency of the IL-36 receptor antagonist) (23, 24). Interestingly, IL-36 γ has been demonstrated to induce prominent production of the potent neutrophil chemoattractant IL-8 (25), which would be compatible with the extensive infiltration of neutrophils seen in skin lesions from patients suffering from acneiform eruptions (5). Furthermore, clinical trial data has shown that subcutaneous anti-IL-8 antibody injection strongly abrogates the induction of acneiform skin toxicity by EGFRi (26). To define the cell types expressing IL-36 γ in the skin of patients with acneiform eruption, immunohistochemical analyses and mRNA in situ hybridization were performed. In line with gene expression data, histochemical analysis of patients' lesions revealed elevated IL-36 γ expression, which was predominantly localized in keratinocytes of epidermal hair follicles (Figure 1C and Supplemental Figure 1B-C). This result and the fact that EGFR is preferentially expressed in undifferentiated and proliferating keratinocytes in the basal and suprabasal layers of the epidermis as well as the outer layers of the hair follicle (5), led to the hypothesis that

keratinocytes might be key players in the acneiform eruption by producing IL-36 γ in response to EGFRi.

EGFRi and C. acnes synergize to promote IL-36 γ expression and skin inflammation

To examine whether EGFR inhibition could lead to enhanced IL-36 γ production in keratinocytes, primary human keratinocytes (PHKs) were exposed to the EGFRi erlotinib in vitro. Upon exposure to 1 μ M erlotinib - a concentration compatible with the serum concentration found in treated patients (27) - PHKs produced 3.2-fold ($p=0.048$) higher levels of IL-36 γ than upon exposure to vehicle alone as quantified by RT-PCR (Figure 1D). Given that both common acne vulgaris and EGFRi-induced eruptions occur in sebum-rich regions of the body that are colonized with *C. acnes*, and that *C. acnes* is known to be involved in the pathogenesis of acne (19-22), we exposed PHKs to both erlotinib and *C. acnes*. Interestingly, IL-36 γ production at the mRNA and protein level was further enhanced (8.4-fold in mRNA, $p=0.001$) when PHKs were simultaneously exposed to erlotinib and *C. acnes* (Figure 1D-E). In contrast, the transcripts of other inflammatory cytokines such as TNF- α , IL-1 β and IL-6 were not significantly increased by simultaneous exposure to erlotinib and *C. acnes* (Figure 1D). Similar levels of IL-36 γ induction in PHKs were also observed with cetuximab, another EGFRi, when used in combination with *C. acnes* (Supplemental Figure 1D). Furthermore, these results were confirmed when EGFR was genetically silenced using siRNA (Supplemental Figure 1E).

Besides IL-36 γ , expression of other genes were significantly increased in acneiform lesions by EGFRi, including the S100 proteins S100A12 and S100A8, the chemokine CXCL6 and the pleiotropic immunomodulatory cytokine IL-24 (Figure 1A). The regulation of the expression of these genes by erlotinib and *C. acnes* was also assessed in PHKs, and with the exception of S100A8, the expression of these genes was not as elevated as that of IL-36 γ (Supplemental Figure 1F). However, the expression of the above transcripts could be significantly induced in PHKs by exposure for 6 hours to IL-36 γ alone (Supplemental Figure 1G), suggesting that IL-36 γ may be an upstream driver cytokine in EGFRi-induced acneiform eruption.

Similar to exposure to the TLR2 agonist *C. acnes* (28-30), IL-36 γ release into the culture supernatant of PHKs could be induced by exposure to erlotinib and the TLR2 agonist Pam3CSK4 (Figure 1F). In line with this, knocking down TLR2 attenuated IL-36 γ production induced by erlotinib and *C. acnes* (Supplemental Figure 1H-I). Expression of the neutrophil chemoattractant IL-8 has previously been shown to be induced by IL-36 γ (23, 25, 31-35) and

was also found by gene expression profiling and quantitative PCR to be upregulated (32-fold, $p=0.041$) in acneiform lesional skin (Figure 1B). To determine if simultaneous EGFR inhibition and TLR2 signaling can trigger IL-36 γ -dependent production of IL-8 in human skin, we exposed normal human skin *ex vivo* to erlotinib and Pam3CSK4. In line with the neutrophil-rich inflammation and enhanced IL-8 gene expression observed in acneiform lesional skin by EGFRi, increased IL-8 production (16-fold, $p=0.0051$) was observed in human skin explants exposed to erlotinib and Pam3CSK4 as compared to vehicle, erlotinib or Pam3CSK4 alone (Figure 1G). In the same *ex vivo* experimental setting, addition of the recombinant IL-36 receptor antagonist to erlotinib and Pam3CSK4 resulted in a significant reduction of IL-8 production (4.1-fold, $p=0.014$) (Figure 1G), thus establishing the IL-36 dependency of IL-8 expression in human skin exposed simultaneously to EGFRi and TLR2 agonists. These data demonstrate that EGFR inhibition and simultaneous TLR2 activation act synergistically to drive keratinocyte IL-36 γ expression with subsequent production of the neutrophil chemoattractant IL-8 in the skin. Taken together with the observed high levels of expression of IL-36 γ and the neutrophil-rich inflammation observed in the pilo-sebaceous units of inflamed skin by EGFRi, this is suggestive of a central pathogenic role of keratinocyte-derived IL-36 γ in the acneiform skin toxicity caused by EGFRi.

Increased expression and binding of the transcription factor KLF4 to the IL-36 γ promoter upon EGFR inhibition

To understand how EGFRi and TLR2 signaling synergistically promote IL-36 γ production in PHKs, we analyzed the transcriptional regulation of human IL-36 γ . Histone modification patterns in PHKs revealed one enhancer and one promoter region upstream of the IL-36 γ gene, the promoter region containing a binding site for the NF- κ B subunit p65 (36) (Supplemental Figure 2A). Interestingly, EGFR inhibition alone, or *C. acnes* exposure alone, resulted in only moderate enhancement of IL-36 γ reporter activity as assessed in a Luciferase reporter assay of human IL-36 γ transcriptional activity in PHKs (Figure 2A). This is suggestive of the existence of two distinct responsive sites in the promoter region of the IL-36 γ gene. In response to EGFR inhibition and *C. acnes* exposure, the activation pattern of a reporter containing both the IL-36 γ enhancer and promoter regions was similar to the pattern observed when only the promoter was present (Supplemental Figure 2B), suggesting that the enhancer region is dispensable for erlotinib and *C. acnes*-induced IL-36 γ production in PHKs. Therefore, and given the synergistic effect of erlotinib and *C. acnes*, this observation suggests that the IL-36 γ promoter contains a binding site for an additional transcription factor to NF- κ B p65. To identify this site, we generated IL-36 γ reporters with promoter deletions of

increasing lengths, thus mapping a genomic region located within 1130 and 1100 bp upstream of the first ATG as crucial for IL-36 γ transcriptional activity (Figure 2B-C). Furthermore, reporters containing mutations in either the genomic region located between 1130 and 1100 bp upstream of the first ATG or within the p65-binding site revealed, respectively, a 42% (p=0.010) and 81% (p=0.0003) reduction in IL-36 γ transcriptional activity, whereas mutation of both regions resulted in a 92% (p=0.0002) reduction (Figure 2D-E), indicating that both regions are required for optimal IL-36 γ transcriptional activity.

To identify the putative transcription factor that binds to the -1130 to -1100 bp region of the IL-36 γ gene promoter, we searched the JASPAR database, an open-access repository for matrix-based transcription factor binding profiles (37), and identified fourteen transcription factors as potential candidates (Supplemental Figure 2C). As no NF- κ B-related transcription factors were revealed by this search, we hypothesized that the EGFRi-responsive site (EiRS) is located in this -1130 to -1100 bp region of the IL-36 γ gene promoter. We subsequently performed quantitative PCR of mRNA derived from keratinocytes exposed to EGFR inhibitor erlotinib, and amongst these 14 candidates thereby identified a significant change in the expression levels of two transcription factors, KLF4 and ZEB1 (Figure 2F). Since ZEB1 transcription decreased after EGFR inhibition, and is weakly expressed in normal keratinocytes (refs. 38, 39 and Supplemental Figure 2D), we considered KLF4 as the probable candidate and assessed whether KLF4 could effectively bind to the EGFRi responsive region of the IL-36 γ promoter. Using an electrophoretic mobility shift assay (EMSA), we could effectively demonstrate that KLF4 specifically binds to the DNA sequence within the -1130 to -1100 bp region of the IL-36 γ gene promoter (Figure 2G). In line with the above, EGFR inhibition resulted in increased KLF4 expression in PHKs (Figure 2H), and DNA pull down assays performed with the same sequence as previously used in the EMSA revealed that KLF4 from erlotinib-exposed PHKs could specifically bind to the EGFR inhibitor-responsive region located between -1130 to -1100 bp upstream of the ATG in the IL-36 γ gene promoter (Figure 2I). In addition, exposure of human skin ex vivo to erlotinib increased the expression of KLF4 (Figure 2J).

Lack of a KLF4 binding site in the mouse IL-36 γ promoter precludes murine EGFRi-induced IL-36 γ response

Next, we examined IL-36 γ production in response to EGFRi and NF- κ B activation in primary murine keratinocytes (PMKs). Surprisingly, despite the ability of the EGFR inhibitor erlotinib to block the phosphorylation of murine EGFR to a similar extent to that of human EGFR (Supplemental Figure 3A), enhanced IL-36 γ production was not observed, in contrast to the

effect observed in human keratinocytes (Figure 3A). In this setting, PMKs were exposed to murine IL-36 γ to achieve NF- κ B activation given their weak response to *C. acnes*, Pam3CSK4 and lipopolysaccharide (LPS) (data not shown). In our culture conditions, PMKs already expressed high levels of KLF4 at the basal state (Supplemental Figure 3B), a characteristic that was irrespective of the numerous culture conditions tested (data not shown). To test the requirement of KLF4 for IL-36 γ transcription in mouse keratinocytes, we compared PMKs from KLF4-knockout mice and wild-type mice, and PMKs overexpressing KLF4, however, without being able to detect synergistic IL-36 γ elevation after NF- κ B activation as observed in human keratinocytes (Figure 3B and Supplemental Figure 3C). In accordance with the above, the putative KLF4 binding site identified by searching the JASPAR database, located at 1140 bp upstream of the first ATG in murine IL-36 γ promoter (Figure 3C), could not be shown by EMSA to form a DNA-protein complex with mouse KLF4 (Figure 3D). Analysis of evolutionarily conserved regions in the genomes of sequenced species revealed that the KLF4 binding region in human IL-36 γ promoter is conserved in the rhesus monkey and chimpanzee, but not in the mouse or rat, whereas the sequence of the IL-36 γ promoter region corresponding to the p65 binding site is approximately 70% conserved in mice and rat as compared to humans (Supplemental Figure 3D and refs. 40, 41). Alignment of the mouse and human IL-36 γ gene loci revealed furthermore that the mouse genome lacks the region corresponding to the 583 bp long-region of human IL-36 γ that contains the KLF4 binding site (-1120 bp) (Supplemental Figure 3E). These results demonstrate that the mouse IL-36 γ promoter is devoid of the KLF4 binding site found in humans, explaining the absence of synergistic induction of IL-36 γ expression by EGFR inhibition and NF- κ B activation in murine keratinocytes, and suggests that the mouse is not an appropriate model for the in vivo analysis of the acneiform skin toxicity to EGFRi.

Blockade of the EGFR-MEK-ERK pathway results in elevated KLF4 and IL-36 γ expression

MEK inhibitors, which block the MAPK-ERK signaling pathway by inhibiting the MAP kinases MEK1 and MEK2, cause adverse skin reactions similar to those observed in EGFRi-treated patients, including the commonly observed acneiform skin toxicity (4). Quantitative PCR analysis of acneiform lesional skin biopsies from MEKi-treated patients revealed, as observed in EGFRi-treated patients, elevated IL-36 γ (9.4-fold, $p=0.0012$) and IL-8 (15-fold, $p=0.019$), but not IL-1 β or IL-6 mRNA levels (Figure 4A and Supplemental Figure 4A). Since MEK is a downstream partner in the EGFR signaling pathway, we next assessed whether MEK inhibition could also result in elevated IL-36 γ gene expression in PHKs. In vitro, the MEK inhibitors trametinib and selumetinib, together with *C. acnes*, synergistically

induced elevated production of IL-36 γ in PHKs, as previously observed with EGFRi (Figure 4B left and Supplemental Figure 4B). Similar results were observed upon ERK silencing with siRNA (Supplemental Figure 4C). Of interest is the reported reduced incidence and severity of cutaneous skin toxicities observed in patients treated simultaneously with a BRAF inhibitor (BRAFi) and MEKi in clinical practice, as compared to patients treated with MEKi alone, and this has been shown to be due to paradoxical ERK activation in BRAF wild-type cells (42-44). In line with this clinical observation, when PHKs were pre-exposed to the BRAFi vemurafenib prior to exposure to trametinib and *C. acnes*, the expression of IL-36 γ mRNA induced by trametinib was significantly inhibited (7.4-fold, p=0.0002) (Figure 4B and Supplemental Figure 4D).

Consistent with increased IL-36 γ expression observed upon inhibition of the EGFR-MEK-ERK pathway at different levels, elevated KLF4 expression was also observed (Figure 4C-D). Furthermore, ERK1 and ERK2 could be co-immunoprecipitated with KLF4 from HEK293T cells transfected with FLAG-tagged KLF4 and Myc-tagged ERK1 and ERK2 (Figure 4E), suggesting possible post-transcriptional modification of KLF4 by ERK1/2. Indeed, enhanced poly-ubiquitination (top panel) and phosphorylation of proline-neighboring serine or threonine residues (second panel) of KLF4 was observed in the presence of constitutively active ERK, and the latter is consistent with the activity of “proline-directed protein kinases” ERKs (45) (Figure 4F). To determine if KLF4 expression is regulated by ubiquitination and proteasomal degradation, expression in response to proteasome inhibition was analyzed. Indeed, increased KLF4 expression was observed upon proteasomal inhibition with MG132 (Supplemental Figure 4E), indicating that KLF4 expression is controlled also post-translationally, and targeted for proteasomal degradation after ERK1/2 phosphorylation, as a downstream consequence of EGFR-MEK pathway activation. These data show that inhibition of either EGFR or MEK signaling in keratinocytes elevates KLF4 expression post-translationally (Supplemental Figure 4F).

KLF4 enhances IL-36 γ transcriptional activity upon EGFR/MEK inhibition

To determine if KLF4 is capable of enhancing IL-36 γ transcriptional activity, we overexpressed KLF4 in PHKs. Such an overexpression resulted in enhanced IL-36 γ expression at the protein level upon exposure of PHKs to *C. acnes* (Figure 5A). Similarly, doxycycline-inducible overexpression of wild-type KLF4 enhanced IL-36 γ transcriptional activity whereas dominant-negative KLF4 mutant did not (Figure 5B). This demonstrates that forced expression of KLF4 can alone mimic the effect of EGFR/MEKi to drive IL-36 γ production in keratinocytes. In accordance with this, siRNA silencing of KLF4 substantially

suppressed the ability of EGFRi and *C. acnes* to enhance IL-36 γ production (Figure 5C). The deletion of KLF4 in keratinocyte cell lines using the CRISPR/Cas9 system resulted in a loss of induction of IL-36 γ gene expression in response to MEKi (Figure 5D and Supplemental Figure 5A). Furthermore, mutation of the KLF4-binding site in keratinocyte cell lines by CRISPR/Cas9 abrogated the ability of MEKi to induce IL-36 γ transcription, whereas in these cell lines IL-1 β expression was unaffected (Figure 5E). This demonstrates an essential role of KLF4 and its binding to the IL-36 γ promoter in regulating IL-36 γ transcriptional activity.

Consistent with a previous report (46), inhibition of the EGFR-MEK-ERK pathway resulted in increased KLF4 expression in the nucleus of PHKs in vitro (Figure 6A). In acneiform skin lesions from EGFRi treated patients, abundant nuclear KLF4 expression could be observed in keratinocytes. In contrast, only low levels of nuclear KLF4 expression was observed in control skin samples (Figure 6B-C). This data suggests that inhibition of either EGFR or MEK signaling enhances nuclear KLF4 expression in keratinocytes in the skin.

Discussion

Here we demonstrate that EGFR and MEK inhibitors partner with the commensal bacterium *C. acnes* that colonizes sebum rich skin to potently induce keratinocyte IL-36 γ expression and drive IL-8-mediated neutrophil rich inflammation, the pathogenic hallmark of the so called “acneiform” skin toxicity frequently associated with these targeted agents. On the basis of in vitro and ex vivo investigations we pinpoint the regulation of keratinocyte IL-36 γ expression upon EGFR blockade to two important signaling events. First, an upregulation of the expression and subsequent binding of the transcription factor KLF4 to its binding site in the promoter region of IL-36 γ , and second, a signal provided by *C. acnes* resulting in the binding of NF- κ B p65 to a sequence in close proximity to the above-mentioned KLF binding site. Interestingly, the simultaneous binding of these two transcription factors to the IL-36 γ promoter results not only in an additive but in a synergistic effect on IL-36 γ gene transcription, which is of relevance for the characteristic localization of the EGFRi induced rash to sebum-rich regions of the skin densely populated with pilo-sebaceous units and *C. acnes* such as the central face, upper chest, and back (17-22). Beyond the topographical distribution of the skin eruption, clinical practice guidelines for therapy support the involvement of *C. acnes* in the pathogenesis of acneiform skin toxicity by EGFR/MEKi. Such guidelines recommend systemic treatment using tetracyclins such as doxycycline or retinoids such as acitretin (47), both of which exert an antibacterial effect on *C. acnes*, either directly for the former, or indirectly by reducing sebum production leading to an alteration of the follicular micromilieu and an indirect reduction in *C. acnes* counts by up to 3 log (48).

As previously reported, the dysfunction of skin barrier and antimicrobial peptide production resulting from EGFR signaling abrogation are important events that can cause severe skin inflammation (13, 14). It is unclear, however, to what extent this may contribute to the initiation of acneiform skin toxicity by EGFR/MEKi, possibly by facilitating the penetration of commensals such as *C. acnes* into the epidermis and/or pilo-sebaceous unit. Indeed, a possible involvement of other stimuli in addition to EGFR inhibition are suggested in skin rash development of a mouse model (49). KLF4 has been demonstrated to be a key driver of terminal epidermal differentiation in the skin (50). The enhanced differentiation induced by increased KLF4 in response to EGFR/MEK inhibition might be an important event leading to the skin barrier dysfunction. KLF4 has only rarely been mentioned to be related to inflammatory diseases, but is known as a regulator of proinflammatory cytokine expression in rheumatoid arthritis (51), and is one of the susceptibility genes for psoriasis (52, 53), two diseases in which IL-36 γ is significantly increased in the inflamed tissues (54-57). Indeed, elevated expression of KLF4 has been reported in synovial tissue from rheumatoid arthritis patients and in the epidermis of psoriatic skin (51, 58), but the exact role of KLF4 in the pathogenesis of these inflammatory diseases remains to be defined.

IL-36 γ has been demonstrated to form a self-amplifying inflammatory loop in keratinocytes which express high levels of the IL-36 receptor (34, 59). Besides DITRA, there is accumulating evidence that IL-36 signaling plays an important role in various neutrophilic dermatoses including generalized pustular psoriasis (GPP), palmo-plantar pustular psoriasis (PPP), acute generalized exanthematous pustulosis (AGEP) and acrodermatitis continua Hallopeau (60, 61). The data presented here provides substantial evidence that acneiform skin toxicity caused by EGFR/MEKi should be added to the growing list of pustular skin diseases in which IL-36 likely plays a central pathogenic role. Our findings provide a basis for understanding the physiopathology of acneiform skin toxicity by caused EGFR/MEKi that may lead to better benefit from the antitumor effects with reduced side effects. Several IL-36 inhibitors have been developed and Phase 2 clinical trials of anti-IL-36 receptor antibody in patients with GPP and PPP are ongoing, suggesting that the latter may also offer a possibility for targeted therapy of acneiform skin toxicities by EGFR/MEKi in the near future.

Materials and Methods

Human skin samples: Biopsies were obtained from lesional skin of EGFR and MEK inhibitor-treated patients with acneiform eruption. Normal skin was obtained from specimens

from the Plastic Surgery Department. All biopsies were immediately frozen in liquid nitrogen and stored at -80°C for RNA extraction or directly fixed in Formalin (4% (g/v)) for at least 24 hours for histology.

Mice: *Klf4*-floxed mice were obtained from MMRRC (Columbia, MO). Rosa26-CreER^{T2} mice were obtained from The Jackson Laboratory (Bar Harbor, ME). Tamoxifen-inducible *Klf4* knockout mice were generated by crossing Rosa26-CreER^{T2} mice and *Klf4*-floxed mice. *Klf4* was knocked out by daily i.p. injection of tamoxifen at a dose of 100 mg/kg for five consecutive days. Wild-type C57BL/6 mice were obtained from Janvier labs (Le Genest-Saint-Isle, France).

Cell culture: Primary human keratinocytes were cultured as previously described (62). Briefly, primary human keratinocytes were isolated from fresh, surgically resected human neonatal foreskin. Keratinocytes were grown in keratinocyte serum free medium (#17005-042, Thermo Scientific, Waltham, MA), supplemented with EGF and BPE (Thermo Scientific) and seeded for experiments after 3 passages. All cells were maintained at 37 °C in a 5% CO₂ humidified atmosphere. Primary mouse keratinocytes were isolated from pooled ear and tail. Briefly, skin specimens were incubated with the dermal side down at 37°C in Dulbecco's modified Eagle's medium (DMEM) supplemented with 1.25% trypsin (Sigma-Aldrich, St. Louis, MI) and antibiotic-antimycotic solution (Gibco BRL, Paisley, Scotland) for 30 minutes. Separated epidermis was minced with sterile scissors and incubated at 37°C in DMEM supplemented with 10% (v/v) fetal bovine serum (FBS) and 0.25 mg/ml DNase I for 30 minutes, followed by filtration through a 70 µm cell strainer (BD Biosciences, San Jose, CA). Cells were resuspended in fresh keratinocyte serum free medium (#10744-019, Thermo Scientific) containing 50 ng/ml EGF (#E4127, Sigma-Aldrich), 10⁻¹⁰ M cholera toxin (#C8052, Sigma-Aldrich) and antibiotic-antimycotic solution (Gibco BRL) and seeded in Corning® Collagen I (15 µg/cm², #354236, Corning, NY)-coated dishes. After one day of attachment, non-adherent cells were washed away and fresh medium was added. HEK293T cells were cultured in DMEM supplemented with 10% (v/v) FBS, antibiotic-antimycotic solution (Gibco BRL), sodium-pyruvate (Invitrogen, Carlsbad, CA) and GlutaMAX solution (Invitrogen). Puromycin (#P9620) and blasticidin (#15205) were from Sigma-Aldrich.

Plasmids: Human genomic DNA was isolated from primary human keratinocytes by QIAamp® DNA Mini Kit (QIAGEN, Venlo, Netherlands). The 6782 bp upstream sequence from the first ATG of human IL-36γ gene was amplified using Pfu polymerase (Invitrogen) with primers (forward; 5'-cacctgggcatattgcataatgg-3', reverse; 5'-aagcttagtggtgtgtctcagcac-3', excluding an additional flanking BglII/HindIII site) and cloned into a luciferase reporter

vector pGL3-Basic (Promega, Fitchburg, WI). The human IL-36 γ promoter (1630 bp) luciferase construct and its NF- κ B mutant construct were gifts from Professor Heiko Mühl (Goethe University Frankfurt, Germany). Site-directed mutagenesis was performed using pfu Turbo (Thermo Scientific) according to manufacturer's instructions to generate a point mutation in the EGFR inhibition-responsive site. Sequentially shorter reporter constructs of the human IL-36 γ promoter were generated from the human IL-36 γ promoter (1630 bp) construct, using following forward primers excluding an additional flanking BglII site; 5'-ccatgtggatggagctgaaa-3' (1180 bp); 5'-gcctggctttccattcaggt-3' (1135 bp); 5'-gtgggtagttgagaaatgc-3' (1105 bp); 5'-cttgctgagacgtgtggct-3' (1076 bp). The dominant-negative human KLF4 construct was generated from human KLF4 construct (Addgene #26815, Cambridge, MA), using a following reverse primer excluding additional restriction enzyme site; 5'-aaagaggggaagacgatcgtaa-3'. The following plasmids were purchased from Addgene and cloned into pcDNA3.1 (Invitrogen) or pMXs-IP (Gift from Prof. Kitamura, University of Tokyo, Japan); mouse Klf4 (Addgene #15920), human ERK1 (#23509), human ERK2 (#23498) and human ubiquitin (#31815). The constitutive active (CA)-ERK plasmid was a gift from Prof. Westermarck (University of Turku, Finland).

Reagents: Erlotinib was purchased from MedChem Express (Monmouth Junction, NJ). Cetuximab was from MERCK Serono (Darmstadt, Germany). Trametinib and vemurafenib were from ApexBio (Boston, MA). Selumetinib and MG132 were from Selleckchem (Houston, TX). Recombinant human IL-36 γ (#6835), mouse IL-36 γ (#6996) and human IL-36Ra (#1275) were from R&D systems (Minneapolis, MN). Pam3CSK4 was from InvivoGen (San Diego, CA). The goat anti-IL-36 γ (#AF2320) and anti-mouse KLF4 (#AF3158) antibodies were from R&D systems (Minneapolis, MN). The rabbit anti-IL-36 γ (#LS-C201142) and its blocking peptide (#LS-E45854) were from LifeSpan BioSciences (Seattle, WA). The anti-human KLF4 antibody (#AM09057PU-N) was from Acris (Germany). The anti- β -actin, anti-FLAG (#F1804) and anti-myc (#C3956) antibodies were from Sigma-Aldrich. The anti-human KLF4 (#12173), anti-ERK (#9107), anti-phospho-ERK (#4370), anti-phospho-threonine/proline (#9391) antibodies were from Cell Signaling Technology (Danvers, MA). The anti-T7 antibody was from Abcam (#ab9138, Cambridge, UK). The anti-HA antibody was from Santa Cruz Biotechnology (#sc-805, Santa Cruz, CA). The secondary antibodies used were alkaline phosphatase-conjugated mouse IgG (#S372B), rabbit IgG (#S373B) and goat IgG (#V115A) from Promega. Live *C. acnes* was prepared as previously described (63).

Gene expression array: Total RNA was extracted from individual skin samples using Tri-Reagent (Sigma-Aldrich) according to manufacturer's instructions. 1 µg of each RNA sample was converted to complementary DNA with RT² First Strand kit (Qiagen) and used in real-time PCR performed on Human Inflammatory Response & Autoimmunity RT² Profiler™ PCR Array (PAHS-3803Z, Qiagen) according to manufacturer's protocol. Data analysis was performed using $\Delta\Delta C_t$ method.

Quantitative PCR: Complementary DNA was generated from total RNA using RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific) according to manufacturer's instructions. Quantitative real-time PCR was performed using a LightCycler®480 (Roche, Basel, Switzerland) with SYBR Green I Master mix (Roche). The primers used for amplification of specific genes were synthesized by Microsynth (Balgach, Switzerland) (**Supplemental Table 2**).

In situ hybridization: 620 nucleotide-long human IL-36 γ cDNA was amplified using Pfu polymerase (Invitrogen) with primers (forward; 5'-ggaagctgctggagccacgattc-3', reverse; 5'-aaagaccaagctgccacctctagg-3', excluding an additional flanking HindIII/EcoRI site) and cloned into pcDNA3 vector (Invitrogen). PCR fragments for probes were amplified with primers for CMV and BGH. The digoxigenin (DIG)-labeled antisense and sense RNA probes for human IL-36 γ were synthesized by in vitro transcription using either SP6 or T7 RNA polymerase with the DIG RNA labeling kit (Roche). These probes were hydrolyzed in hydrolysis buffer (40mM NaHCO₃, 60mM Na₂CO₃) to be 0.25 kb. The unincorporated nucleotides were removed using a spin column (Roche). Formalin-fixed paraffin-embedded tissue sections (4 µm) were deparaffinized and rehydrated in RNase-free condition. Sections were treated with 1 µg/ml proteinase K for 20 minutes and washed with 2xSSC for three times, followed by prehybridization for 2 h in 2xSSC containing 50 % formamide. Hybridization buffer (HB) contained 50% formamide, 4xSSC, 100 ng/ml yeast tRNA and 10% dextran sulfate. 25 ng DIG-labeled RNA probes were diluted in 50 µl HB, heated to 95 °C for 5 minutes, added to the tissues and hybridized overnight. After hybridization, the tissues were washed and incubated in stringent wash buffer (20% formamide, 2xSSC) at 42 °C for 30 minutes, followed by 2 µg/mL RNase A treatment at 37 °C for 1 hour. The sections were washed in 2xSSC and 0.2xSSC at 55°C each for 30 minutes and PBS at RT for 5 minutes. Blocking was performed in 5% BSA in PBS at RT for 1 h and sections were incubated in AP-conjugated anti-DIG Fab fragment (1/4000 dilution; Roche) at RT for 2 hours. After three washes in PBS for 5 minutes, tissues were stained using 2% NBT/BCIP in 0.1M NaCl, 0.1M Tris-HCl (pH 9) at RT in the dark for two days.

Immunohistochemistry: 5 μ m formalin-fixed, paraffin-embedded human skin sections were deparaffinized and rehydrated. Antigen unmasking was performed by heating the slides for 25 minutes in Target Retrieval solution (DAKO, Glostrup, Denmark). Sections were blocked using 5% BSA in PBS for 1 hour and stained for 2 hours at room temperature with anti-IL-36 γ antibody. Primary antibodies were detected using a biotin-conjugated secondary antibody (Southern Biotech, Birmingham, USA) followed by an Avidin-Biotin-complex and addition of peroxidase substrate (Vector Laboratories, Peterborough, UK). Nuclei were counterstained using a solution of hematoxylin. The sections were mounted in mounting medium (DAKO, Santa Clara, CA) and imaged using an Aperio ScanScope (Leica Biosystems, Wetzlar, Germany).

Immunofluorescence staining: PHKs were seeded on circular 18-mm glass coverslips (Hecht-Assisten, Sondheim/Rhön, Germany). Cells were fixed for 30 minutes in 3% paraformaldehyde/2% sucrose solution, permeabilized for 2 minutes with 0.2% Triton X-100 in phosphate buffered saline and blocked for 1 hour in 1% bovine serum albumin (BSA Fraction V; GE Healthcare) in 0.5% Tween-20 in PBS. In vitro keratinocyte samples were stained for 2 hours at room temperature with anti-KLF4 goat antibody followed by DyLight488-conjugated secondary antibody (ab96891; Abcam, Cambridge, MA). Skin tissue samples were prepared as described in the immunohistochemistry paragraph. The sections were stained for 2 hours at room temperature with anti-KLF4 mouse antibody and anti-IL-36 γ rabbit antibody followed by Alexa Fluor 488/555-conjugated secondary antibodies (A-11001, A-21429; Thermo Scientific). Nuclei were counterstained with DAPI. The sections were mounted in the mounting medium and imaged using an Aperio ScanScope (Leica Biosystems, Wetzlar, Germany).

Western blot: To prepare whole cell lysates, cells were lysed in SDS buffer with DTT (Sigma-Aldrich). Proteins were separated on SDS-PAGE gels with Mini-PROTEAN Tetra Vertical Electrophoresis Cell (Bio-Rad Laboratories, Inc., Hercules, CA) at a constant voltage (80-120V) and transferred to an Amersham Protran 0.2 μ m nitrocellulose membrane (GE Healthcare) using semi-dry or wet systems from Bio-Rad. The membranes were blocked with 5% dried milk in PBS supplemented with 0.5% Tween-20 (Sigma-Aldrich) and then probed overnight with primary antibodies at 4 $^{\circ}$ C followed by AP-conjugated secondary antibodies for 1 hour at room temperature. Proteins were detected using BCIP/NBT color development substrate (Promega) and dried membranes were scanned using a LiDE210 scanner (Canon Inc., Tokyo, Japan).

ChIP-seq/DNase-seq/RNA-seq data analysis: Raw sequencing data were converted to Fastq files by NCBI SRA Toolkit. Quality control on the raw data was performed by FastQC. The reads for ChIP-seq/DNase-seq were aligned to human reference genome (build GRCh37/hg19) by Bowtie2 Aligner. The mapped sequence reads were transformed to a binary format, sorted and indexed by SAMtools, followed by generation of coverage plots by BEDtools. These files were converted into BigWig files by BedGraphToBigWig and visualized in IGV. ChIP-seq data from primary human keratinocytes were obtained from the following ChIP-seq/DNase-seq samples: GSM941735, GSM733698, GSM733674, GSM733636 and GSM816635. ChIP-seq data of p65 was from GSM935526. The reads for RNA-seq were aligned to human reference genome (build GRCh37/hg19) by HISAT2. The transcripts were assembled by Cufflinks, followed by generation of differential gene expression data by Cuffdiff. RNA-seq data from primary human keratinocytes were obtained from GSM2074746, GSM2074747 and GSM2074748.

Luciferase reporter assay: Human and mouse primary keratinocytes were transfected with reporter constructs using TransIT-X2® Dynamic Delivery System (Mirus Bio, Madison, WI) according to manufacturer's instructions. Co-transfection of the Renilla-luciferase expression vector pRL-TK (Promega, Fitchburg, WI) was used as an internal control for all reporter assays. Cell extracts were generated 24 hours post transfection using Reporter Lysis Buffer (Promega) and extracts were assayed for firefly luciferase and Renilla-luciferase activity using the Luciferase Assay system (Promega) and coelenterazine (0.1 µg/ml, Sigma-Aldrich), respectively. Luminescence was measured with the Cytation3 Imaging Reader (BioTek, Winooski, VT).

Gene Transfer and knockdown: HEK293T cells were transfected with 8 µg of mammalian expression construct of human and mouse KLF4 using TransIT-X2® Dynamic Delivery System (Mirus Bio). Primary human keratinocytes were transduced using a published protocol (64) with minor modifications. Briefly, viral supernatant was produced by transfecting 8 µg of KLF4-pMXs-IP to Phoenix Ampho cells (ATCC, Manassas, VA) using TransIT-X2® Dynamic Delivery System (Mirus Bio). Viral supernatant was collected 48 hours after transfection and added to keratinocytes in 6-well plates supplemented with 10 µg/mL polybrene (Sigma-Aldrich) followed by centrifugation at 650 g for 45 minutes at 32°C. After centrifugation, keratinocytes were washed with PBS and cultured in fresh medium. The same transfection step was repeated in the next day and keratinocytes were incubated another 24 hours for experiments. Mouse keratinocytes were transfected in 12-well plates with 3 µg of mammalian expression construct of mouse KLF4 using the TransIT-X2®

Dynamic Delivery System (Mirus Bio). Silencing RNA (siRNA) transfection of primary human keratinocytes was carried out using INTERFERin (Polyplus-transfection, Illkirch, France) at a final concentration of 5 nM endoribonuclease-prepared siRNA (esiRNA) according to manufacturer's protocol. The esiRNAs used (EGFR, KLF4, ERK1, ERK2) were purchased from Sigma-Aldrich. Experiments were performed 2 days after transfection. EGFP esiRNA was used as a control. Short hairpin RNA (shRNA) fragments of human TLR2 were hybridized with synthesized sense and anti-sense oligonucleotides. The sense strand sequence is 5'-ccggccagccagaaagcactacaatctcgagattgtagtgctttctggctggtttt-3'. DNA oligonucleotides were synthesized by Microsynth and ligated into Tet-pLKO-puro (Addgene #21915). Viral supernatant was produced by transfecting the Tet-pLKO-puro plasmid, psPAX2 (Addgene, #12260) and pMD2.G (Addgene, #12259) into HEK 293T cells and viruses were harvested 48 hours later. KERTr keratinocyte cell lines (ATCC CRL-1658, Manassas, VA) were transduced with the virus and selected for 1 week with 1 µg/ml puromycin. Gene knockdown was induced by culturing the cells in the medium containing 1 µg/ml doxycycline.

Generation of CRISPR/Cas9-cell lines: Single-stranded DNA oligonucleotides (Supplemental Table S3) were cloned into pLentiCRISPRv2 plasmid (Addgene, #98293). Viral supernatant was produced by transfecting the pLentiCRISPRv2 plasmid, psPAX2 and pMD2.G into HEK 293T cells and viruses were harvested 48 hours later. KERTr keratinocyte cell lines were transduced with the virus and selected for 10 days with 10 µg/ml blasticidin. Cloning was performed by limiting dilution in conditioned medium. Genomic DNA was extracted from the isolated single cell-expanded clones using QIAamp DNA Mini Kit and amplicons harboring the targeted alleles were prepared by PCR using Taq polymerase (#EP0404, Thermo Scientific). The PCR amplicons were cloned into a TOPO vector using TA Cloning Kit (Promega) according to the manufacturer's protocol and then sent for Sanger sequencing.

EMSA: The sequences of the probes used for EMSA were; EGFR inhibition-responsive site wild-type forward; 5'-ttcattcaggtgtggccttag-3', wild-type reverse; 5'-ctaagccacac~~ct~~gaatggaa-3', mutant forward; 5'-ttcattca~~aat~~gtggccttag-3', mutant reverse; 5'-ctaagccacatttgaatggaa-3', putative murine KLF4 binding site forward; 5'-gagatccaggtggaaggaaga-3', and reverse; 5'-tcttcttccacctggatctc-3'. The probes including Cy3-modification at the 5' end were synthesized by Microsynth. To construct oligonucleotide duplexes, 2 nmol of each sense and antisense oligonucleotides were annealed in a buffer (100 mM NaCl, 50 mM Tris-HCl, 10 mM MgCl₂ and 100 µg/ml BSA) by heating the mixtures to 95°C for 5 minutes and allowing the solution to cool slowly to room temperature. EMSA was

performed with 5 µg of cell lysate, 0.15 pmol of Cy3-labeled oligonucleotides, 1.5 µg of BSA, 0.5 µg of poly(dI-dC) and in 12 µl of reaction mixture (24 mM HEPES-KOH pH 7.9, 8 mM Tris-HCl (pH 8.0), 2 mM EDTA, 1 mM DTT and 12% glycerol) with a proteinase inhibitor cocktail (Roche). Competition assays were performed to demonstrate the sequence specific binding of the probes. For the competition assays, a 50-fold molar excess of unlabeled wild-type or mutant oligonucleotide probe was added 20 minutes before the addition of Cy3-labeled probes and incubated for another 20 minutes at room temperature. Supershift assays were performed to demonstrate the complex formation of the protein of interest and the target probe, by means of appearance of a new supershifted band upon addition of antibody targeting the protein of interest. For the supershift assays, 1 µg of anti-KLF4 antibody (Cell Signaling Technology) was added to the reaction mixture for 20 minutes before the addition of Cy3-labeled probes. Samples were loaded onto Novex® 6% DNA retardation gels (Thermo Scientific) and subjected to electrophoresis in 0.5X Tris borate buffer for 30 minutes at 150 mV. Gels were subsequently visualized on an Odyssey® Fc Imaging System (LI-COR Biosciences, France).

DNA pull down assay: The sequences of the probes used for the DNA pull down assay were the same as those for EMSA. 5'-biotinylated wild-type and mutant forward oligonucleotides were synthesized by Microsynth and annealed with non-biotinylated reverse oligonucleotides. Cell lysates were incubated in the same reaction buffer as EMSA with 7.5 nmol of double-stranded annealed oligonucleotides at 4°C overnight with gentle shaking on a rocker. 30 µl of Pierce™ High Capacity Streptavidin Agarose beads (Thermo Scientific) was added and incubated at 4°C for 1 hour. Beads were washed five times with ice-cold PBS using SigmaPrep™ spin column (Sigma-Aldrich). Proteins bound to streptavidin beads were dissolved in 2x SDS sample buffer, boiled at 95 °C for 5 minutes and subjected to immunoblotting.

Co-immunoprecipitation analysis: HEK293T cells were transfected in 6-well plates with 2.4 µg of plasmids (Myc-tagged ERK1 (0.4 µg) and ERK2 (2 µg) and FLAG-tagged KLF4 (0.4 µg); HA-tagged ubiquitin (0.8 µg), FLAG-tagged KLF4 (0.8 µg) and CA-ERK (0.8 µg)) using TransIT-X2® Dynamic Delivery System (Mirus) according to manufacturer's instructions. Empty vector pcDNA3.1 was added to be 2.4 µg of plasmids in total. 24 hours after transfection, cells were washed with ice-cold PBS and lysed with pre-chilled lysis buffer (50 mM Tris-Hcl (pH 7.5), 100 mM NaCl and 0.1% (v/v) Triton-X supplemented with protease and phosphatase inhibitor cocktail (Roche) for 30 minutes on a rocker at 4 °C. Samples for ubiquitin analysis were boiled at 95°C for 5 minutes in 1% SDS and sample

buffer was added to dilute to 0.1% SDS. Samples were incubated overnight with anti-FLAG (1 µg) at 4 °C, followed by the addition of 30 µl protein A/G PLUS Agarose beads (Santa Cruz Biotechnology) and incubation for 2 hours. Immunoprecipitates were collected by centrifugation at 1,000 g for 5 minutes at 4 °C and the beads were then washed five times with lysis buffer. The beads were resuspended in 2x SDS sample buffer, boiled at 95 °C for 5 minutes and subjected to immunoblotting.

ELISA: Primary human keratinocytes were exposed to 1 µM erlotinib and 5 µg/mL Pam3CSK4 in 6-well plates for 48 hours. After washing three times with pre-warmed PBS, cells were incubated in fresh medium for 48 hours. After centrifugation, supernatants were collected and subjected to IL-36γ ELISA (Adipogen, Lausanne, Switzerland).

Ex vivo skin culture: Ex vivo skin culture was performed using a published protocol (65) with the following modifications. Full-thickness skin specimens were obtained from patients undergoing plastic or reconstructive surgery. Skin samples were cut to small pieces (4 x 4 mm) and placed in 24-well plates containing 0.5 ml of keratinocyte serum free medium (#17005-042, Thermo Scientific) supplemented with EGF, BPE (Thermo Scientific), 100 µg/mL kanamycin (Invitrogen) and 1.4 mM CaCl₂ (Sigma-Aldrich). Tissue cultures were then incubated at 37 °C in a 5% CO₂ atmosphere with fresh culture medium provided at 2 day intervals. Ex vivo skin explants were cultured with 1 µg/mL erlotinib for 24 hours for western blotting or for 4 days followed by snap freezing for subjecting to quantitative PCR.

Statistical analysis: Statistical analysis was performed using unpaired Student's t-test or one-way analysis of variance (ANOVA) followed by Dunnett's multiple-comparison test, using Prism 7.02 software (GraphPad, La Jolla, CA). Differences were considered significant when: *P ≤ 0.05, **P ≤ 0.01 and ***P ≤ 0.001.

Study approval: All experiments with human samples and the use of human skin samples for research studies were carried out in accordance with the Cantonal Ethical Committee of Zurich, Switzerland after informed written patient consent and according to the Declaration of Helsinki Principles. All animal procedures were approved by the Cantonal Veterinary Office of Zurich, Switzerland.

Acknowledgments: We thank R. Dummer, S. Goldinger, G. Restivo and D. Hug for collecting skins from patients and healthy individuals. The authors thank S. Grossi, M.C. Brüggen, D. Mohanan, R. Huber, B. Kiefer, I.S. Kleiber, A. Kolios, M. Yokota, N. Li, A. Klug, B. Schmid, P. Bosshard, J. Danis and M. Tusup for technical assistance. This work was supported by grants from the Swiss National Science Foundation (grant 310030-176035 to

L.E.F.) and the Zurich University Research Priority Program “Translational Cancer Research” to L.E.F. M.M. acknowledges support from radiz-Rare Disease Initiative Zürich, Clinical Research Priority Program for Rare Diseases University of Zürich, the Monique Dornonville de la Cour Foundation and the Novartis Foundation for Medical-Biological Research.

Author contributions: T.K.S., E.C. and L.E.F. designed the research; T.K.S., E.C. and L.E.F. supervised the research; T.K.S., M.M, G.F. and A.O. conducted the studies; B.MS., HD.B, T.R., JT.M., J.H. and A.A.N. collected human biological samples and patients’ data. T.K.S., M.M., E.C. and L.E.F. analysed the data; T.K.S., M.M., E.C. and L.E.F. prepared the manuscript. T.K.S., M.M., E.C. and L.E.F. revised the manuscript. All authors reviewed the manuscript.

Competing Financial Interests: The authors declare no competing financial interests.

References

1. Sibilila M, Kroismayr R, Lichtenberger BM, Natarajan A, Hecking M, Holcman M. The epidermal growth factor receptor: from development to tumorigenesis. *Differentiation*. 2007;75(9):770-787.
2. Ciardiello F, Tortora G. EGFR antagonists in cancer treatment. *N Engl J Med*. 2008;358(11):1160-1174.
3. Caunt CJ, Sale MJ, Smith PD, Cook SJ. MEK1 and MEK2 inhibitors and cancer therapy: the long and winding road. *Nat Rev Cancer*. 2015;15(10):577-592.
4. Zhao Y, Adjei AA. The clinical development of MEK inhibitors. *Nat Rev Clin Oncol*. 2014;11(7):385-400.
5. Lacouture ME. Mechanisms of cutaneous toxicities to EGFR inhibitors. *Nat Rev Cancer*. 2006;6(10):803-812.
6. Lacouture ME, Balagula Y. Acneiform eruption secondary to epidermal growth factor receptor (EGFR) inhibitors. In: Mockenhaupt M, ed. *UpToDate*. Waltham, MA.: UpToDate; 2017. www.uptodate.com. Accessed November 30, 2017.
7. Boone SL, Rademaker A, Liu D, Pfeiffer C, Mauro DJ, Lacouture ME. Impact and management of skin toxicity associated with anti-epidermal growth factor receptor therapy: survey results. *Oncology*. 2007;72(3-4):152-159.
8. Osio A, Mateus C, Soria JC, et al. Cutaneous side-effects in patients on long-term treatment with epidermal growth factor receptor inhibitors. *Br J Dermatol*. 2009;161(3):515-521.
9. Wu PA, Balagula Y, Lacouture ME, Anadkat MJ. Prophylaxis and treatment of dermatologic adverse events from epidermal growth factor receptor inhibitors. *Curr Opin Oncol*. 2011;23(4):343-351.
10. Wacker B, Nagrani T, Weinberg J, Witt K, Clark G, Cagnoni PJ. Correlation between development of rash and efficacy in patients treated with the epidermal growth factor receptor tyrosine kinase inhibitor erlotinib in two large phase III studies. *Clin Cancer Res*. 2007;13(13):3913-3921.
11. Stintzing S, Kapaun C, Laubender RP, et al. Prognostic value of cetuximab-related skin toxicity in metastatic colorectal cancer patients and its correlation with parameters of the epidermal growth factor receptor signal transduction pathway: results from a randomized trial of the GERMAN AIO CRC Study Group. *Int J Cancer*. 2013;132(1):236-245.
12. Liu HB, Wu Y, Lv TF, et al. Skin rash could predict the response to EGFR tyrosine kinase inhibitor and the prognosis for patients with non-small cell lung cancer: a systematic review and meta-analysis. *PLoS one*. 2013;8(1):e55128.
13. Mascia F, Lam G, Keith C, et al. Genetic ablation of epidermal EGFR reveals the dynamic origin of adverse effects of anti-EGFR therapy. *Sci Transl Med*. 2013;5(199):199ra110.
14. Lichtenberger BM, Gerber PA, Holcman M, et al. Epidermal EGFR controls cutaneous host defense and prevents inflammation. *Sci Transl Med*. 2013;5(199):199ra111.
15. Franzke CW, Cobzaru C, Triantafyllidou A, et al. Epidermal ADAM17 maintains the skin barrier by regulating EGFR ligand-dependent terminal keratinocyte differentiation. *J Exp Med*. 2012;209(6):1105-1119.
16. Yang J, Meyer M, Muller AK, et al. Fibroblast growth factor receptors 1 and 2 in keratinocytes control the epidermal barrier and cutaneous homeostasis. *J Cell Biol*. 2010;188(6):935-952.

17. Bossi P, Liberatoscioli C, Bergamini C, et al. Previously irradiated areas spared from skin toxicity induced by cetuximab in six patients: implications for the administration of EGFR inhibitors in previously irradiated patients. *Ann Oncol.* 2007;18(3):601-602.
18. Mitra SS, Simcock R. Erlotinib induced skin rash spares skin in previous radiotherapy field. *J Clin Oncol.* 2006;24(16):e28-29.
19. Grice EA, Kong HH, Conlan S, et al. Topographical and temporal diversity of the human skin microbiome. *Science.* 2009;324(5931):1190-1192.
20. O'Neill AM, Gallo RL. Host-microbiome interactions and recent progress into understanding the biology of acne vulgaris. *Microbiome.* 2018;6(1):177.
21. Hall JB, Cong Z, Imamura-Kawawasa Y, et al. Isolation and Identification of the Follicular Microbiome: Implications for Acne Research. *J Invest Dermatol.* 2018;138(9):2033-2040.
22. Fitz-Gibbon S, Tomida S, Chiu BH, et al. Propionibacterium acnes strain populations in the human skin microbiome associated with acne. *J Invest Dermatol.* 2013;133(9):2152-2160.
23. Marrakchi S, Guigue P, Renshaw BR, et al. Interleukin-36-receptor antagonist deficiency and generalized pustular psoriasis. *N Engl J Med.* 2011;365(7):620-628.
24. Onoufriadis A, Simpson MA, Pink AE, et al. Mutations in IL36RN/IL1F5 are associated with the severe episodic inflammatory skin disease known as generalized pustular psoriasis. *Am J Hum Genet.* 2011;89(3):432-437.
25. Towne JE, Garka KE, Renshaw BR, Virca GD, Sims JE. Interleukin (IL)-1F6, IL-1F8, and IL-1F9 signal through IL-1Rrp2 and IL-1RAcP to activate the pathway leading to NF-kappaB and MAPKs. *J Biol Chem.* 2004;279(14):13677-13688.
26. Bangsgaard N, Houtkamp M, Schuurhuis DH, et al. Neutralization of IL-8 prevents the induction of dermatologic adverse events associated with the inhibition of epidermal growth factor receptor. *PLoS one.* 2012;7(6):e39706.
27. Hidalgo M, Siu LL, Nemunaitis J, et al. Phase I and pharmacologic study of OSI-774, an epidermal growth factor receptor tyrosine kinase inhibitor, in patients with advanced solid malignancies. *J Clin Oncol.* 2001;19(13):3267-3279.
28. Kim J, Ochoa MT, Krutzik SR, et al. Activation of toll-like receptor 2 in acne triggers inflammatory cytokine responses. *J Immunol.* 2002;169(3):1535-1541.
29. Nagy I, Pivarcsi A, Koreck A, Szell M, Urban E, Kemeny L. Distinct strains of Propionibacterium acnes induce selective human beta-defensin-2 and interleukin-8 expression in human keratinocytes through toll-like receptors. *J Invest Dermatol.* 2005;124(5):931-938.
30. Lheure C, Grange PA, Ollagnier G, et al. TLR-2 Recognizes Propionibacterium acnes CAMP Factor 1 from Highly Inflammatory Strains. *PLoS one.* 2016;11(11):e0167237.
31. Foster AM, Baliwag J, Chen CS, et al. IL-36 promotes myeloid cell infiltration, activation, and inflammatory activity in skin. *J Immunol.* 2014;192(12):6053-6061.
32. Towne JE, Renshaw BR, Douangpanya J, et al. Interleukin-36 (IL-36) ligands require processing for full agonist (IL-36alpha, IL-36beta, and IL-36gamma) or antagonist (IL-36Ra) activity. *J Biol Chem.* 2011;286(49):42594-42602.
33. Li N, Yamasaki K, Saito R, et al. Alarmin function of cathelicidin antimicrobial peptide LL37 through IL-36gamma induction in human epidermal keratinocytes. *J Immunol.* 2014;193(10):5140-5148.
34. Friedrich M, Tillack C, Wollenberg A, Schaubert J, Brand S. IL-36gamma sustains a proinflammatory self-amplifying loop with IL-17C in anti-TNF-induced psoriasiform skin lesions of patients with Crohn's disease. *Inflamm Bowel Dis.* 2014;20(11):1891-1901.
35. Chustz RT, Nagarkar DR, Poposki JA, et al. Regulation and function of the IL-1 family cytokine IL-1F9 in human bronchial epithelial cells. *Am J Respir Cell Mol Biol.* 2011;45(1):145-153.
36. Bachmann M, Scheiermann P, Hardle L, Pfeilschifter J, Muhl H. IL-36gamma/IL-1F9, an innate T-bet target in myeloid cells. *J Biol Chem.* 2012;287(50):41684-41696.
37. Khan A, Fornes O, Stigliani A, et al. JASPAR 2018: update of the open-access database of transcription factor binding profiles and its web framework. *Nucleic Acids Res.* 2018;46(D1):D260-D266.
38. Mehrzarin S, Chen W, Oh JE, et al. The p63 Gene Is Regulated by Grainyhead-like 2 (GRHL2) through Reciprocal Feedback and Determines the Epithelial Phenotype in Human Keratinocytes. *J Biol Chem.* 2015;290(32):19999-20008.
39. Lee B, Villarreal-Ponce A, Fallahi M, et al. Transcriptional mechanisms link epithelial plasticity to adhesion and differentiation of epidermal progenitor cells. *Dev Cell.* 2014;29(1):47-58.
40. Ovcharenko I, Nobrega MA, Loots GG, Stubbs L. ECR Browser: a tool for visualizing and accessing data from comparisons of multiple vertebrate genomes. *Nucleic Acids Res.* 2004;32(Web Server issue):W280-286.
41. Aken BL, Ayling S, Barrell D, et al. The Ensembl gene annotation system. *Database (Oxford).* 2016;2016:baw093.
42. Robert C, Karaszewska B, Schachter J, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med.* 2015;372(1):30-39.

43. Flaherty KT, Robert C, Hersey P, et al. Improved survival with MEK inhibition in BRAF-mutated melanoma. *N Engl J Med.* 2012;367(2):107-114.
44. Gibney GT, Messina JL, Fedorenko IV, Sondak VK, Smalley KS. Paradoxical oncogenesis--the long-term effects of BRAF inhibition in melanoma. *Nat Rev Clin Oncol.* 2013;10(7):390-399.
45. Cano E, Mahadevan LC. Parallel signal processing among mammalian MAPKs. *Trends Biochem Sci.* 1995;20(3):117-122.
46. Kim MO, Kim SH, Cho YY, et al. ERK1 and ERK2 regulate embryonic stem cell self-renewal through phosphorylation of Klf4. *Nat Struct Mol Biol.* 2012;19(3):283-290.
47. Lacouture ME, Anadkat MJ, Bensadoun RJ, et al. Clinical practice guidelines for the prevention and treatment of EGFR inhibitor-associated dermatologic toxicities. *Support Care Cancer.* 2011;19(8):1079-1095.
48. King K, Jones DH, Daltrey DC, Cunliffe WJ. A double-blind study of the effects of 13-cis-retinoic acid on acne, sebum excretion rate and microbial population. *Br J Dermatol.* 1982;107(5):583-590.
49. Surguladze D, Deevi D, Claros N, et al. Tumor necrosis factor-alpha and interleukin-1 antagonists alleviate inflammatory skin changes associated with epidermal growth factor receptor antibody therapy in mice. *Cancer Res.* 2009;69(14):5643-5647.
50. Sen GL, Boxer LD, Webster DE, et al. ZNF750 is a p63 target gene that induces KLF4 to drive terminal epidermal differentiation. *Dev Cell.* 2012;22(3):669-677.
51. Luo X, Chen J, Ruan J, et al. Kruppel-Like Factor 4 Is a Regulator of Proinflammatory Signaling in Fibroblast-Like Synoviocytes through Increased IL-6 Expression. *Mediators Inflamm.* 2016;2016:1062586.
52. Tsoi LC, Spain SL, Knight J, et al. Identification of 15 new psoriasis susceptibility loci highlights the role of innate immunity. *Nat Genet.* 2012;44(12):1341-1348.
53. Madonna S, Scarponi C, Sestito R, Pallotta S, Cavani A, Albanesi C. The IFN-gamma-dependent suppressor of cytokine signaling 1 promoter activity is positively regulated by IFN regulatory factor-1 and Sp1 but repressed by growth factor independence-1b and Kruppel-like factor-4, and it is dysregulated in psoriatic keratinocytes. *J Immunol.* 2010;185(4):2467-2481.
54. Boutet MA, Bart G, Penhoat M, et al. Distinct expression of interleukin (IL)-36alpha, beta and gamma, their antagonist IL-36Ra and IL-38 in psoriasis, rheumatoid arthritis and Crohn's disease. *Clin Exp Immunol.* 2016;184(2):159-173.
55. Mahil SK, Catapano M, Di Meglio P, et al. An analysis of IL-36 signature genes and individuals with IL1RL2 knockout mutations validates IL-36 as a psoriasis therapeutic target. *Sci Transl Med.* 2017;9(411).
56. Keermann M, Koks S, Reimann E, Prans E, Abram K, Kingo K. Transcriptional landscape of psoriasis identifies the involvement of IL36 and IL36RN. *BMC Genomics.* 2015;16:322.
57. Johnston A, Xing X, Guzman AM, et al. IL-1F5, -F6, -F8, and -F9: a novel IL-1 family signaling system that is active in psoriasis and promotes keratinocyte antimicrobial peptide expression. *J Immunol.* 2011;186(4):2613-2622.
58. Kim KJ, Park S, Park YH, et al. The expression and role of kruppel-like factor 4 in psoriasis. *Ann Dermatol.* 2014;26(6):675-680.
59. Carrier Y, Ma HL, Ramon HE, et al. Inter-regulation of Th17 cytokines and the IL-36 cytokines in vitro and in vivo: implications in psoriasis pathogenesis. *J Invest Dermatol.* 2011;131(12):2428-2437.
60. Satoh TK, Mellett M, Contassot E, French LE. Are neutrophilic dermatoses autoinflammatory disorders? *Br J Dermatol.* 2016;178(3):603-613.
61. Meier-Schiesser B, Feldmeyer L, Jankovic D, et al. Culprit Drugs Induce Specific IL-36 Overexpression in Acute Generalized Exanthematous Pustulosis. *J Invest Dermatol.* 2019;139(4):848-858.
62. Feldmeyer L, Keller M, Niklaus G, Hohl D, Werner S, Beer HD. The inflammasome mediates UVB-induced activation and secretion of interleukin-1beta by keratinocytes. *Curr Biol.* 2007;17(13):1140-1145.
63. Kistowska M, Gehrke S, Jankovic D, et al. IL-1beta drives inflammatory responses to propionibacterium acnes in vitro and in vivo. *J Invest Dermatol.* 2014;134(3):677-685.
64. Aasen T, Izpisua Belmonte JC. Isolation and cultivation of human keratinocytes from skin or plucked hair for the generation of induced pluripotent stem cells. *Nat Protoc.* 2010;5(2):371-382.
65. Tavakkol A, Varani J, Elder JT, Zouboulis CC. Maintenance of human skin in organ culture: role for insulin-like growth factor-1 receptor and epidermal growth factor receptor. *Arch Dermatol Res.* 1999;291(12):643-651.

Figure 1




Figure 1: Increased production of IL-36 γ in primary keratinocytes and lesional skin of patients suffering from acneiform eruptions in response to EGFR inhibition and *C. acnes*. (A) Gene expression profiling from lesional skin of five patients and five healthy control (HC). Heatmap of the top 12 most differentially expressed genes ranked from lowest false discovery rate (FDR) and 12 selected genes are shown. (B) Quantitative PCR of mRNA from lesional skin samples of 10 EGFR inhibitor-treated patients with acneiform eruption and 10 healthy control skin biopsies. Data represent mean (SD). (C) Immunohistochemistry staining with goat anti-IL-36 γ antibody of formalin-fixed paraffin-embedded skin sections of acneiform eruption patient and normal donors. Scale bar represents 100 μ m. Pictures are representative of five patients and five healthy individuals. (D) PHKs were exposed to erlotinib (EGFR inhibitor, 1 μ M) and *C. acnes* (MOI of 10) for 6 hours. Total RNA was subjected to qPCR analysis. Data represent means \pm SEM. $n = 3$. (E) PHKs were exposed to either erlotinib (1 μ M) or *C. acnes* (MOI of 10) or both for 24 hours. Cell lysates were subjected to western blotting using specific antibodies to IL-36 γ and β -actin. Blots were run contemporaneously with the same protein samples. (F) PHKs were exposed to erlotinib (1 μ M) and Pam3CSK4 (5 μ g/mL). IL-36 γ secretion was measured by ELISA in culture supernatants. Data represent means \pm SEM. $n = 3$. (G) Ex vivo skin explants were exposed to erlotinib (1 μ M), Pam3CSK4 (5 μ g/mL) and/or human IL-36Ra (1 μ g/mL). The skin samples were then subjected to quantitative PCR. Data represent means \pm SEM. $n = 4$. Data were analyzed with 2-tailed unpaired t test (B), 1-way ANOVA followed by Dunnett's (D and F) or Tukey's multiple-comparisons test (G). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Data are representative of 3 independent experiments.

Figure 2




Figure 2. KLF4 binds to the IL-36 γ promoter and regulates IL-36 γ transcriptional activity in response to EGFR inhibition. (A) Luciferase reporter assay of human IL-36 γ transcriptional activity in PHKs transfected with IL-36 γ -pGL3 (1630 bp) reporter plasmid, followed by exposure to erlotinib and *C. acnes* for 16 hours. TK Renilla luciferase was measured to determine transfection efficiency. Data represent means \pm SEM. *n*=4. (B, D) Schematic of 5'-deletion and mutant constructs of the human IL-36 γ promoter. Site-directed mutagenesis was performed to introduce the indicated mutation at the EGFR inhibitor-responsive site (EiRS) and p65 binding site. (C, E) 5'-Deletion and mutation study of the human IL-36 γ promoter activity. PHKs were transfected with indicated plasmids, followed by exposure to erlotinib and *C. acnes* for 16 hours. *n*=4, Data represent means \pm SEM. (F) RT-PCR was performed to evaluate the gene expression of transcription factor candidates binding to the EiRS. PHKs were exposed to erlotinib for 6 hours. Data represent means \pm SEM. *n*=3. (G) Gel shift, competition and supershift EMSA analysis using Cy5-labeled oligonucleotide probe of the EiRS-containing region and HEK293T cell lysate containing KLF4 protein. (H) PHKs were exposed to erlotinib and *C. acnes* for 24 hours. (I) DNA pull-down assay using biotinylated wild-type- or mutant-oligonucleotide probe of the EiRS-containing region. These probes were incubated with extracts from PHKs exposed to erlotinib for 24 hours. DNA-associated proteins were visualized by western blotting. (J) Ex vivo skin explants from healthy controls were exposed to erlotinib for 24 hours and KLF4 expression was assessed by western blotting. The blot shown is representative of two different skin donors. Data were analyzed with 1-way ANOVA followed by Dunnett's multiple-comparisons test (A and E) or with 2-tailed unpaired Mann-Whitney *U* (C) or t test (F). **P*<0.05, ***P*<0.01, ****P*<0.001. All blots were run contemporaneously with the same protein samples.

Figure 3




Figure 3. Lack of KLF4 binding site results in loss of synergistic IL-36 γ production in mice. (A) PMKs were exposed to erlotinib (1 μ M) and murine IL-36 γ (100 ng/mL) for 6 hours, isolated RNA was subject to qPCR. Data represent means \pm SEM. $n = 3$. (B) PMKs from wild-type or KLF4 knockout mouse were exposed to murine IL-36 γ (100 ng/mL) for 6 hours. Data represent means \pm SEM. $n = 3$. PMK cell lysates were subject to SDS-PAGE and immunoblotting. Blots were run contemporaneously with the same protein samples. (C) Schematic of the human and murine IL-36 γ promoter with predicted KLF4 binding site and p65 binding site by JASPAR. (D) Gel shift and competition EMSA analysis using Cy5-labeled oligonucleotide probe of human and mouse KLF4 binding sites and HEK293T cell lysate containing murine KLF4 protein. Sequence specific binding of human probe to murine KLF4 was demonstrated as a positive control. Gel shift reflecting formation of protein-DNA complexes with the murine probe and KLF4 was not observed. Data are representative of 3 independent experiments.

Figure 4




Figure 4: Blockade of the EGFR-MEK-ERK pathway increases keratinocyte expression of KLF4. (A) Quantitative PCR was performed to evaluate gene expression in RNA isolated from biopsies of 4 patients with acneiform eruption and 10 healthy control skin biopsies. Data represent mean (SD). (B) PHKs were pre-exposed to the BRAF inhibitor vemurafenib (1 $\mu\text{g}/\text{mL}$) for 30 minutes and exposed to the MEK inhibitor trametinib (2 $\mu\text{g}/\text{mL}$) and *C. acnes* (MOI of 10) for 6 hours. Data represent means \pm SEM. $n = 3$. Data were analyzed with 2-tailed unpaired t test (A) or 1-way ANOVA followed by Tukey's multiple-comparisons test (B). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. (C) PHKs were exposed to erlotinib (1 μM) or trametinib (2 $\mu\text{g}/\text{mL}$) for 24 hours and total cell lysates were collected for western blotting with antibodies to KLF4 and β -actin. (D) ERK1 and ERK2 were silenced by siRNA in PHKs and cell lysates were subjected to SDS-PAGE and immunoblotting with indicated antibodies. (E) HEK293T cells were transfected with FLAG-tagged KLF4 and Myc-tagged ERK1 and ERK2 for 24 hours. Cell lysates were immunoprecipitated with an anti-FLAG antibody, followed by immunoblotting with the indicated antibodies. (F) HEK293T cells were transfected with FLAG-tagged KLF4, HA-tagged ubiquitin and constitutively active ERK for 24 hours. Cell lysates were immunoprecipitated with an anti-FLAG antibody, followed by immunoblotting with the indicated antibodies. All blots were run contemporaneously with the same protein samples. Data are representative of 3 independent experiments.

Figure 5




Figure 5: KLF4 is critical for IL-36 γ transcriptional activity upon EGFR/MEK inhibition. (A) KLF4 overexpressing primary keratinocytes were exposed to *C. acnes* for 24 hours. (B) Flag-tagged wild-type (WT)- and dominant-negative (DN)-KLF4 were overexpressed in response to doxycycline using Tet-on system for 24 hours, followed by exposure to Pam3CSK4 for another 24 hours. The cell lysates were collected for western blotting and qPCR. Data represent means \pm SEM. $n = 3$. (C) KLF4 siRNA-treated PHKs were exposed to erlotinib and *C. acnes* for 6 hours. Data represent means \pm SEM. $n = 3$. (D) KLF4 knocked out keratinocyte cell lines by CRISPR/Cas9 were exposed to trametinib (2 μ g/mL) for 24 hours and total cell lysates were collected for western blotting with antibodies to KLF4 and β -actin. The cells were exposed to trametinib for 24 hours and isolated RNA was subject to qPCR. Data represent means \pm SEM. $n = 3$. All blots were run contemporaneously with the same protein samples. Data are representative of 3 independent experiments. (E) Mutations generated by CRISPR/Cas9 in the KLF4 binding site. Red nucleotides are the PAM sequence and blue nucleotides hybridize to the sgRNA. KLF4 binding site-mutant cells were exposed to trametinib and Pam3CSK4 for 24 hours. Total RNA was subjected to qPCR analysis. Data represent means \pm SEM. $n = 3$. Data were analyzed with 1-way ANOVA followed by Dunnett's (B and E) or Tukey's multiple-comparisons test (C) or with 2-tailed unpaired t test (D). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure 6




Figure 6: Increased KLF4 level in vitro and in vivo upon EGFR inhibition. (A)

Representative images of KLF4 expression (green) in PHKs after erlotinib or control DMSO exposure for 24 hours. Nuclei were stained with DAPI. Scale bar represents 10 μm . Data are representative of 3 independent experiments. **(B-C)** Immunofluorescence staining with mouse anti-KLF4 (green) and rabbit anti-IL-36 γ (red) antibodies of formalin-fixed paraffin-embedded skin sections of acneiform eruption patients and normal donors. Nuclei were stained with DAPI. The white box regions were zoomed separately in the figure C. Scale bar represents 50 μm **(B)** and 10 μm **(C)**. Pictures are representative of five patients and three healthy individuals.