

PKC λ regulates glucose-induced insulin secretion through modulation of gene expression in pancreatic β cells

Naoko Hashimoto, ... , Tetsuo Noda, Masato Kasuga

J Clin Invest. 2005;115(1):138-145. <https://doi.org/10.1172/JCI22232>.

Article

Metabolism

Altered regulation of insulin secretion by glucose is characteristic of individuals with type 2 diabetes mellitus, although the mechanisms that underlie this change remain unclear. We have now generated mice that lack the λ isoform of PKC in pancreatic β cells (β PKC $\lambda^{-/-}$ mice) and show that these animals manifest impaired glucose tolerance and hypoinsulinemia. Furthermore, insulin secretion in response to high concentrations of glucose was impaired, whereas the basal rate of insulin release was increased, in islets isolated from β PKC $\lambda^{-/-}$ mice. Neither the β cell mass nor the islet insulin content of β PKC $\lambda^{-/-}$ mice differed from that of control mice, however. The abundance of mRNAs for Glut2 and HNF3 β was reduced in islets of β PKC $\lambda^{-/-}$ mice, and the expression of genes regulated by HNF3 β was also affected (that of *Sur1* and *Kir6.2* genes was reduced, whereas that of *hexokinase 1* and *hexokinase 2* genes was increased).

Normalization of HNF3 β expression by infection of islets from β PKC $\lambda^{-/-}$ mice with an adenoviral vector significantly reversed the defect in glucose-stimulated insulin secretion. These results indicate that PKC λ plays a prominent role in regulation of glucose-induced insulin secretion by modulating the expression of genes important for β cell function.

Find the latest version:

<https://jci.me/22232/pdf>





PKC λ regulates glucose-induced insulin secretion through modulation of gene expression in pancreatic β cells

Naoko Hashimoto,¹ Yoshiaki Kido,¹ Tohru Uchida,¹ Tomokazu Matsuda,¹ Kazuhisa Suzuki,² Hiroshi Inoue,¹ Michihiro Matsumoto,¹ Wataru Ogawa,¹ Sakan Maeda,³ Hiroaki Fujihara,⁴ Yoichi Ueta,⁴ Yasuo Uchiyama,⁵ Kazunori Akimoto,⁶ Shigeo Ohno,⁶ Tetsuo Noda,⁷ and Masato Kasuga¹

¹Department of Clinical Molecular Medicine, Division of Diabetes and Digestive and Kidney Diseases, ²Department of Development and Aging, Division of Internal and Geriatric Medicine, and ³Department of Biomedical Informatics, Division of Molecular Pathology, Kobe University Graduate School of Medicine, Kobe, Japan. ⁴Department of Physiology, University of Occupational and Environmental Health, School of Medicine, Kitakyushu, Japan. ⁵Department of Cell Biology and Neuroscience, Osaka University School of Medicine, Osaka, Japan. ⁶Department of Molecular Biology, Yokohama City University Graduate School of Medical Science, Yokohama, Japan. ⁷Department of Experimental Pathology, Cancer Institute, Tokyo, Japan.

Altered regulation of insulin secretion by glucose is characteristic of individuals with type 2 diabetes mellitus, although the mechanisms that underlie this change remain unclear. We have now generated mice that lack the λ isoform of PKC in pancreatic β cells (β PKC $\lambda^{-/-}$ mice) and show that these animals manifest impaired glucose tolerance and hypoinsulinemia. Furthermore, insulin secretion in response to high concentrations of glucose was impaired, whereas the basal rate of insulin release was increased, in islets isolated from β PKC $\lambda^{-/-}$ mice. Neither the β cell mass nor the islet insulin content of β PKC $\lambda^{-/-}$ mice differed from that of control mice, however. The abundance of mRNAs for Glut2 and HNF3 β was reduced in islets of β PKC $\lambda^{-/-}$ mice, and the expression of genes regulated by HNF3 β was also affected (that of *Sur1* and *Kir6.2* genes was reduced, whereas that of *hexokinase 1* and *hexokinase 2* genes was increased). Normalization of HNF3 β expression by infection of islets from β PKC $\lambda^{-/-}$ mice with an adenoviral vector significantly reversed the defect in glucose-stimulated insulin secretion. These results indicate that PKC λ plays a prominent role in regulation of glucose-induced insulin secretion by modulating the expression of genes important for β cell function.

Introduction

Type 2 diabetes mellitus is characterized by insulin resistance in peripheral tissues and functional failure of pancreatic β cells. The importance of the β cells of islets of Langerhans in the pathogenesis of type 2 diabetes is reflected in the development of potential new treatments such as islet transplantation and regenerative islet cell therapy. The roles of molecules important in β cell function in vivo have recently begun to be examined by gene targeting. Signaling by receptor tyrosine kinases has thus been implicated in the regulation both of β cell mass (1, 2) and of insulin secretion (3–5). For example, deletion of the insulin receptor specifically in mouse β cells resulted in impairment of glucose-stimulated insulin secretion and a reduction in β cell mass with age, effects that led to the development of glucose intolerance (6). Ablation of the IGF-1 receptor in mouse β cells also resulted in impaired insulin secretion but did not affect β cell morphology (7, 8). Silencing of the insulin receptor or IGF-1 receptor by RNA interference in MIN6 cells blocked activation of PI3K and consequently inhibited glucose-induced insulin secretion (9). 3-Phosphoinositide-dependent kinase-1 is thought to be a key mediator of PI3K signaling and contributes to the activation of AGC protein kinases, including Akt, p70 S6 kinase, and atypical isoforms of PKC (10). Mice that express a constitutively active form of Akt1 exhibit an increased

β cell mass (11, 12), whereas ablation of p70 S6 kinase 1 was associated with a reduced β cell size (13). Atypical PKC isoforms also might therefore be expected to participate in regulation of β cell growth and insulin secretion. In fact, the atypical isoform PKC λ is expressed in pancreatic β cells (14). There have been so far several reports suggesting the role of atypical PKC in insulin secretion (15, 16), insulin synthesis (15, 16), and proliferation (17) in pancreatic β cells. However, direct evidence for the function of PKC λ in β cells in vivo has not been available.

The purpose of this study was to establish a mouse line in which the PKC λ gene is deleted specifically in pancreatic β cells in order to determine the role of this isozyme in these cells in vivo. Our results show that glucose-stimulated insulin secretion is altered in such mice in a manner similar to that apparent in type 2 diabetes, and they suggest that this change is attributable to altered regulation of gene expression important for β cell function.

Results

Generation of β cell-specific PKC λ knockout mice. Pancreatic β cell-specific PKC λ knockout (β PKC $\lambda^{-/-}$) mice were generated by breeding of mice (PKC $\lambda^{lox/lox}$) that harbor a modified endogenous PKC λ gene, in which exon 5 is flanked by loxP sites (K. Akimoto et al., unpublished observations), with mice that express Cre recombinase under the control of the promoter of the rat insulin 2 gene (RIP-Cre mice) (18). To evaluate the efficiency of Cre recombinase expression in pancreatic β cells, we performed PCR analysis with primers targeted to regions external to the loxP sites

Conflict of interest: The authors have declared that no conflict of interest exists.

Citation for this article: *J. Clin. Invest.* 115:138–145 (2005).
doi:10.1172/JCI200522232.




Figure 1

Generation of β cell-specific PKC λ knockout mice. (A) PCR analysis of genomic DNA isolated from islets of control ($PKC\lambda^{flox/flox}$) and $\beta PKC\lambda^{-/-}$ mice. The primers were targeted to regions external to the loxP sites of $PKC\lambda^{flox/flox}$ mice. (B and C) Immunoblot analysis of PKC λ in the islets (B) or the brain, hypothalamus, heart, liver, skeletal muscle, and fat (C) of control and $\beta PKC\lambda^{-/-}$ mice. Tissue homogenates were subjected to immunoprecipitation and subsequent immunoblot analysis with antibodies against PKC λ . The expression of PKC λ in islets was similarly analyzed. (D) Immunostaining of islets in pancreatic sections of control and $\beta PKC\lambda^{-/-}$ mice with antibodies against PKC λ and FITC-conjugated secondary antibodies. Scale bars: 50 μ m.

of $PKC\lambda^{flox/flox}$ mice and with genomic DNA extracted from the islets of $\beta PKC\lambda^{-/-}$ mice and $PKC\lambda^{flox/flox}$ mice (used as control animals thereafter). Whereas the control animals yielded a PCR product of about 1,600 bp, the $\beta PKC\lambda^{-/-}$ mice yielded a major product of about 200 bp and a minor product of about 1,600 bp (Figure 1A), which indicates that the loxP-flanked region containing exon 5 of the PKC λ gene was efficiently removed by Cre recombinase in the β cells of $\beta PKC\lambda^{-/-}$ mice.

We next examined the expression of PKC λ in various tissues by immunoblot analysis. The amount of PKC λ in islets of $\beta PKC\lambda^{-/-}$ mice was reduced by about 80% compared with that in islets of control mice (Figure 1B), indicating the virtually complete loss of PKC λ expression in β cells, given that these cells account for about 80% of the islet cell mass. In contrast, the expression of PKC ζ , another atypical isoform of PKC, was not affected in the islets of $\beta PKC\lambda^{-/-}$ mice (Figure 1B). The abundance of PKC λ in the brain, hypothalamus,

heart, liver, skeletal muscle, and fat was also similar for $\beta PKC\lambda^{-/-}$ and control mice (Figure 1C). Furthermore, hypothalamus from $\beta PKC\lambda^{-/-}$ and control mice was hybridized in situ to probe the complementarity to loxP-flanked exon 5 of the PKC λ gene. The expression of PKC λ mRNA was marginally reduced by about 20% in $\beta PKC\lambda^{-/-}$ mice (data not shown). Immunostaining of pancreatic sections confirmed the efficient ablation of PKC λ in β cells of $\beta PKC\lambda^{-/-}$ mice (Figure 1D). Together, these results indicate that loss of PKC λ expression occurred mainly in pancreatic β cells of $\beta PKC\lambda^{-/-}$ mice.

Metabolic characteristics of $\beta PKC\lambda^{-/-}$ mice. The rate of increase in body weight did not differ between $\beta PKC\lambda^{-/-}$ and control mice (Figure 2A). To assess the possible effects of PKC λ ablation in β cells on glucose metabolism, we measured blood glucose and plasma insulin concentrations in 2- and 6-month-old $\beta PKC\lambda^{-/-}$ and control mice. There were no significant differences in blood glucose and plasma insulin levels between $\beta PKC\lambda^{-/-}$ and control mice at 2 months. At 6 months, the blood glucose concentration of $\beta PKC\lambda^{-/-}$ mice was significantly lower than that of control mice in the fasting state (Figure 2B). The blood glucose level in the fed state as well as plasma insulin concentrations in the fasting or fed state did not differ between the 2 groups of animals, however. Intraperitoneal glucose




Figure 2

Effect of β cell-specific ablation of PKC λ on glucose metabolism. (A) Growth curves of control (open circles) and $\beta PKC\lambda^{-/-}$ (filled circles) mice. Mice were weighed at 2, 4, and 6 months. (B) Blood glucose and plasma insulin concentrations of 6-month-old control (white bars) and $\beta PKC\lambda^{-/-}$ (black bars) mice in the fasting or fed state. (C) Intraperitoneal glucose tolerance tests performed in control and $\beta PKC\lambda^{-/-}$ mice that had fasted overnight. (D) Blood glucose concentrations during insulin tolerance testing in control and $\beta PKC\lambda^{-/-}$ mice. Data are means \pm SE of values from 25 (A), 28 (B), 10 (C), or 6 (D) animals of each genotype. * $P < 0.05$ (ANOVA) versus the corresponding value for control mice.




Figure 3

Impairment of glucose-stimulated insulin secretion in isolated islets of β PKC $\lambda^{-/-}$ mice. Insulin release in response to the indicated concentrations of glucose (A), KCl (B), or tolbutamide (C) was measured with islets isolated from control (white bars) or β PKC $\lambda^{-/-}$ (black bars) mice. Data are means \pm SE of values from 6 animals of each genotype. * P < 0.05 (ANOVA) versus the corresponding value for control mice.

islets at high glucose concentrations. The islets of β PKC $\lambda^{-/-}$ mice exhibited a normal secretory response to a high concentration of KCl (Figure 3B), which elicits insulin release by inducing membrane depolarization. These results thus indicate that depletion of PKC λ in β cells results in a specific impairment in glucose-stimulated insulin secretion and consequent glucose intolerance.

Morphology and insulin content of islets of β PKC $\lambda^{-/-}$ mice. We next determined the possible effects of β cell-specific deficiency of PKC λ on pancreatic morphology. Immunostaining of pancreatic sections from 6-month-old animals with antibodies against insulin and against glucagon revealed a normal islet architecture in β PKC $\lambda^{-/-}$ mice (Figure 4A). Quantitative analysis also revealed no significant difference in the β cell area per pancreas between β PKC $\lambda^{-/-}$ and control mice (Figure 4B). The α cell area per pancreas was also normal in β PKC $\lambda^{-/-}$ mice (data not shown). Furthermore, the total insulin content of islets of β PKC $\lambda^{-/-}$ mice was similar to that of control islets (Figure 4C). These results indicate that islet growth and insulin biosynthesis are not impaired in β PKC $\lambda^{-/-}$ mice. Electron microscopy also revealed that the ultrastructure of β cells, including the number and distribution of insulin-containing dense-core granules, is similar in β PKC $\lambda^{-/-}$ and control mice (Figure 4D), which suggests that the impairment of glucose-stimulated insulin secretion in β PKC $\lambda^{-/-}$ mice is not due

tolerance tests in animals that had fasted overnight revealed that β PKC $\lambda^{-/-}$ mice had abnormal glucose tolerance, with the blood glucose concentration 60 minutes after the glucose load being significantly higher in these mice than in the control animals (Figure 2C). The insulin response to glucose was also impaired in β PKC $\lambda^{-/-}$ mice compared with that in control mice, with the plasma insulin level 60 minutes after the glucose load being significantly lower in the former than in the latter. Insulin tolerance tests did not reveal a significant difference in the insulin sensitivity of peripheral tissues between control and β PKC $\lambda^{-/-}$ mice (Figure 2D), which excludes the possibility that β PKC $\lambda^{-/-}$ mice are insulin resistant.

To investigate further the effect of PKC λ ablation on insulin secretion, we examined glucose- or KCl-stimulated insulin release with islets isolated from β PKC $\lambda^{-/-}$ and control mice. Glucose induced a concentration-dependent increase in insulin secretion from control islets in static incubations, with the extent of insulin release at 16.8 mM glucose being about 20 times that at 2.8 mM glucose (Figure 3A). In contrast, the insulin response of islets from β PKC $\lambda^{-/-}$ mice was significantly greater (3-fold at 2.8 mM glucose) than that of control islets at low glucose concentrations but significantly smaller (about 50% at 16.8 mM glucose) than that of control





Figure 4

Lack of effect of β cell-specific ablation of PKC λ on islet morphology and insulin content. (A) Immunostaining of pancreatic sections from 6-month-old control and β PKC $\lambda^{-/-}$ mice with antibodies against insulin (red) and glucagon (green). Scale bars: 50 μ m. (B) Quantitation of β cell area as a percentage of total pancreatic area in control and β PKC $\lambda^{-/-}$ mice. Data are means \pm SE of values from 4 mice of each genotype. (C) Insulin content of isolated islets. Data are means \pm SE of values from 4 mice of each genotype. (D) Electron microscopy of β cells of control and β PKC $\lambda^{-/-}$ mice. Magnification, \times 10,000.

**Figure 5**

Effects of a high-fat diet on β cell phenotype in $\beta PKC\lambda^{-/-}$ mice. **(A)** Blood glucose and plasma insulin concentrations in the fed state of control and $\beta PKC\lambda^{-/-}$ mice on a high-fat diet. Data are means \pm SE of values from 12 animals of each genotype. **(B)** Intraperitoneal glucose tolerance tests in control (open circles) and $\beta PKC\lambda^{-/-}$ (filled circles) mice after 15 weeks on the high-fat diet. Data are means \pm SE of values from 6 animals of each genotype. * $P < 0.05$ (ANOVA) versus the corresponding value for control mice. **(C)** Immunostaining of pancreatic sections from control and $\beta PKC\lambda^{-/-}$ mice after 15 weeks on the high-fat diet with antibodies against insulin (red) and glucagon (green). Scale bars: 100 μ m. **(D)** Quantitation of β cell area as a percentage of total pancreatic area in control and $\beta PKC\lambda^{-/-}$ mice after 15 weeks on the high-fat diet. Data are means \pm SE of values from 3 mice of each genotype.

to an obvious disruption of cellular architecture or a failure to form insulin secretory granules.


Effect of a high-fat diet on insulin secretion in $\beta PKC\lambda^{-/-}$ mice. We investigated the effect of PKC λ deficiency on β cell function in the insulin-resistant state by feeding control and $\beta PKC\lambda^{-/-}$ mice a high-fat diet for up to 21 weeks from weaning. The increases in body weight were similar for the 2 types of mice on this diet (data not shown). Random measurement of blood glucose concentration in the fed state also did not reveal any significant difference between $\beta PKC\lambda^{-/-}$ and control mice (Figure 5A). The plasma insulin concentration in the fed state tended to be lower in $\beta PKC\lambda^{-/-}$ mice than in control animals, although this difference was not statistically significant. The blood glucose concentration apparent 60 minutes after a glucose challenge was significantly higher in $\beta PKC\lambda^{-/-}$ mice than in controls (Figure 5B). The plasma insulin concentrations apparent after a glucose load were higher in control mice fed the high-fat diet than in those fed a normal diet, reflecting the insulin-resistant state of the former. The insulin levels after a glucose challenge were markedly lower in $\beta PKC\lambda^{-/-}$ mice fed the high-fat diet than in control mice on this diet.

We also examined islet morphology in mice fed the high-fat diet. Compared with those fed normal chow, control mice fed the high-fat diet exhibited an increase in islet mass to compensate for their insulin resistance. Islet mass in $\beta PKC\lambda^{-/-}$ mice fed the high-fat diet was similar to that in the control animals on this diet (Figure 5, C and D). These results indicate that the β cells of $\beta PKC\lambda^{-/-}$ mice grow like those of control mice during the development of insulin resistance, whereas the insulin secretory response to glucose is blunted in the knockout mice.

Expression of genes important for β cell function in $\beta PKC\lambda^{-/-}$ mice. We analyzed the expression of several genes important for β cell function by real-time RT-PCR analysis of total RNA isolated from islets (Figure 6). The amounts of mRNAs for the glucose transporter Glut2 and the Sur1 and Kir6.2 subunits of the ATP-sensitive K⁺ channel

were significantly reduced by 25–35% in $\beta PKC\lambda^{-/-}$ mice compared with those in control mice, which suggests that the impairment of glucose-stimulated insulin secretion in the former animals might result, at least in part, from a deficiency of glucose-sensing proteins. In accordance, the islets of $\beta PKC\lambda^{-/-}$ mice exhibited reduced secretory response to tolbutamide, a sulfonylurea reagent, which elicits insulin release by the closure of the ATP-sensitive K⁺ channel (Figure 3C). The abundance of the mRNA for the transcription factor HNF3 β was also reduced by 45% in the islets of $\beta PKC\lambda^{-/-}$ mice. In contrast, the amounts of mRNAs for the high-affinity hexokinase isoforms 1 and 2 were increased in $\beta PKC\lambda^{-/-}$ mice. The abundance of mRNAs for glucokinase, Pdx1, HNF1 α , HNF1 β , and HNF4 α did not differ between $\beta PKC\lambda^{-/-}$ and control mice. Gene expression profiling with an oligonucleotide microarray confirmed the results obtained by real-time RT-PCR analysis (data not shown).

Restoration of PKC λ expression in islets of $\beta PKC\lambda^{-/-}$ mice. We next examined the effect of restoration of PKC λ expression with an adenoviral vector in islets from $\beta PKC\lambda^{-/-}$ mice on glucose-stimulated insulin secretion. Transfection of islets from control or $\beta PKC\lambda^{-/-}$ mice with a control vector encoding β -galactosidase (AxCALacZ) had no effect on glucose-induced insulin release (data not shown).

**Figure 6**

Altered gene expression in the islets of $\beta PKC\lambda^{-/-}$ mice. The abundance of mRNAs for the indicated proteins was determined by real-time RT-PCR analysis of total RNA isolated from islets of control and $\beta PKC\lambda^{-/-}$ mice. The amounts of the mRNAs in $\beta PKC\lambda^{-/-}$ mice are expressed relative to those in control animals. Data are means \pm SE of triplicates for pooled total RNA samples from 6 mice of each genotype and are representative of a total of 3 similar experiments. * $P < 0.05$ (ANOVA) versus the corresponding value (1.0) for control mice.



The abundance of PKC λ in β PKC $\lambda^{-/-}$ mouse islets transfected with the vector for wild-type PKC λ (AxCA λ wt) was approximately twice that of endogenous PKC λ in islets from control mice (Figure 7A). Restoration of PKC λ expression in islets from the knockout mice normalized the insulin secretory response to glucose (Figure 7B); the hypersecretion of insulin at low glucose concentrations and the impaired insulin secretion at high glucose concentrations apparent with β PKC $\lambda^{-/-}$ mouse islets were thus both significantly reversed by expression of recombinant PKC λ .

The expression of *HNF3 β* and *Kir6.2* genes was also increased in β PKC $\lambda^{-/-}$ mouse islets transfected with AxCA λ wt compared with that in those transfected with AxCALacZ (Figure 7C). Similarly, the increased abundance of *hexokinase 1* mRNA apparent in β PKC $\lambda^{-/-}$ mouse islets was reduced by restoration of PKC λ expression. These results thus indicate that the altered patterns of both insulin secretion and islet gene expression apparent in β PKC $\lambda^{-/-}$ mice are directly attributable to the lack of PKC λ .

Effect of HNF3 β on glucose-stimulated insulin secretion in β PKC $\lambda^{-/-}$ mouse islets. Our results suggested that HNF3 β might be a key mediator of PKC λ action in the regulation of insulin secretion in β cells. We therefore examined the effect of expression of recombinant wild-type HNF3 β in islets isolated from β PKC $\lambda^{-/-}$ mice on glucose-stimulated insulin secretion. Transfection of β PKC $\lambda^{-/-}$ mouse islets with an adenoviral vector for HNF3 β (AxCAHNF3 β) restored the abundance of this transcription factor to the level apparent in islets from control mice (Figure 7D). The insulin secretory response of β PKC $\lambda^{-/-}$ mouse islets to a high glucose concentration was also significantly increased by expression of recombinant HNF3 β (Figure 7E). The increase in the basal secretion of insulin by β PKC $\lambda^{-/-}$ mouse islets also appeared to be partially reversed by transfection with AxCAHNF3 β , although this effect was not statistically significant. These results indicate that HNF3 β might function downstream of PKC λ in the regulation of glucose-induced insulin secretion.

Discussion

We have shown that mice deficient in PKC λ in pancreatic β cells exhibit an increased basal rate of insulin secretion and impairment of glucose-stimulated insulin secretion without any obvious abnormality in β cell morphology. Both glucose-induced transcriptional activation of the insulin gene promoter and glucose-stimulated insulin secretion were previously shown to be inhibited by the PKC-specific inhibitor calphostin C (Ro31-8220) but not by Go6976, an inhibitor of classical and novel PKC isoforms; this implicated atypical PKC in these effects of glucose (15, 16). With the use of isoform-specific pseudosubstrates to inhibit either classical (α , β , and γ) isoforms of PKC or the atypical isoform PKC ζ , Buteau et al. (17) showed that the latter enzyme mediates glucagon-like peptide-1-induced proliferation of pancreatic β cells. However, most of the previous studies of the possible roles of atypical isoforms of PKC in β cells have been performed with inhibitors, and the antibody used to detect atypical PKC did not discriminate between the λ and ζ isoforms. To obtain direct evidence for

the function of PKC λ in β cells, we therefore generated mice that lack this isozyme specifically in these cells with the use of the Cre-loxP system. No compensatory increase in the abundance of PKC ζ was apparent in the islets of the β PKC $\lambda^{-/-}$ mice. RIP-Cre mice have been demonstrated to express Cre recombinase in pancreatic β cells, and also in a subset of neurons in hypothalamus (19). In situ hybridization of brain slices for PKC λ revealed a marginal decrease (approximately 20%) in β PKC $\lambda^{-/-}$ mice. It is possible that the reduced expression of PKC λ in hypothalamus might affect insulin secretory response via the neuronal pathway. However, we believe that the phenotype of β PKC $\lambda^{-/-}$ mice is mainly attributable to the decrease in PKC λ levels in pancreatic β cells, because (a) we observed the impaired glucose-induced insulin secretion and the altered gene expression in islets isolated from β PKC $\lambda^{-/-}$ mice *ex vivo*, and (b) *ex vivo* normalization of PKC λ expression in islets isolated from β PKC $\lambda^{-/-}$ mice resulted in a significant restoration




Figure 7

Effects of adenovirus-mediated restoration of PKC λ or HNF3 β expression on insulin secretion in islets of β PKC $\lambda^{-/-}$ mice. (A–C) Islets isolated from control mice or β PKC $\lambda^{-/-}$ mice were infected with an adenovirus encoding either β -galactosidase (AxCALacZ) or wild-type PKC λ (AxCA λ wt). The islets were then either subjected to immunoblot analysis with antibodies against PKC λ or β actin (A); assayed for insulin secretion in the presence of 2.8 or 16.8 mM glucose (white bars, control islets plus AxCALacZ; black bars, β PKC $\lambda^{-/-}$ islets plus AxCALacZ; gray bars, β PKC $\lambda^{-/-}$ islets plus AxCA λ wt) (B); or subjected to real-time RT-PCR analysis of mRNAs for HNF3 β , hexokinase 1, or Kir6.2 (C). (D and E) Islets isolated from control or β PKC $\lambda^{-/-}$ mice were infected with either AxCALacZ or an adenovirus encoding wild-type HNF3 β (AxCAHNF3 β). The islets were then either subjected to immunoblot analysis with antibodies against HNF3 β or β actin (D) or assayed for insulin secretion in the presence of 2.8 or 16.8 mM glucose (E). Data are means \pm SE of values from 6 mice (B and E) or of triplicates for pooled total RNA samples from 5 mice (C). * P < 0.05 (ANOVA) for the indicated comparisons.



of the impairment of insulin secretion and an alteration of gene expression (Figure 7, A–C).

Complexes of atypical PKC isoforms, PAR-3, and PAR-6 have been shown to be indispensable for cell polarization in *Caenorhabditis elegans* embryos (20), *Drosophila* epithelial cells (21) and neuroblasts (22), and mammalian epithelial cells (23). Expression of a dominant negative mutant of PKC λ thus disrupted the formation of apical-basal polarity in cultured Madin-Darby canine kidney cells (23). We therefore examined whether ablation of PKC λ resulted in disruption of β cell polarity and dysregulation of insulin secretion in β PKC $\lambda^{-/-}$ mice. Electron microscopy did not reveal any obvious structural abnormality related to β cell polarity in β PKC $\lambda^{-/-}$ mice, however. In addition, β PKC $\lambda^{-/-}$ mice manifested impairment of insulin secretion stimulated by glucose but not of that stimulated by KCl, suggesting that the exocytosis of insulin granules remained intact. Together, these results indicate that PKC λ does not contribute substantially to the polarity or structure of β cells, at least in our mice, in which the PKC λ gene is deleted in mature β cells after the onset of Cre expression driven by the rat insulin 2 gene promoter. Several previous studies have also suggested that PKC λ mediates growth factor signaling that leads to cell proliferation, differentiation, and survival (24–26). However, islet size and insulin content did not differ substantially between β PKC $\lambda^{-/-}$ and control mice. Moreover, the extent of β cell apoptosis appeared unaffected in β PKC $\lambda^{-/-}$ mice (data not shown).

PKC λ mediates induction of the expression of the gene for SREBP-1c by either insulin or active PI3K in hepatocytes (27). Indeed, PKC λ contains a functional nuclear localization signal in its NH₂-terminal region (28) and undergoes rapid nucleocytoplasmic shuttling in response to stimuli such as PDGF and nerve growth factor (29). Atypical PKC has also been implicated in the phosphorylation and activation of the transcription factor Pdx1 in response to glucose stimulation in β cells (15). IGF-1 increased the transcriptional activity of the gene for the cytochrome P450 side-chain cleavage enzyme by promoting an interaction between PKC λ and polypyrimidine tract-binding protein-associated splicing factor in the nucleus, and it increased the amount of PKC λ in the nucleus of a granulosa cell line (30). These various observations suggest that PKC λ participates in nuclear events such as the regulation of gene expression.

With the use of quantitative gene expression analysis, we have now shown that ablation of PKC λ affected the expression of genes that encode components of the glucose-sensing mechanism of β cells. The lack of PKC λ thus resulted in increased expression of the genes for the high-affinity hexokinase isoforms 1 and 2, which mediate glucose sensing in the physiological range of blood glucose concentration. Overexpression of hexokinase 1 or 2 has previously been shown to result in an increase in basal insulin release from β cells (31, 32), similar to that observed in the present study with islets derived from β PKC $\lambda^{-/-}$ mice. The loss of PKC λ from β cells also resulted in downregulation of the expression of the genes for Sur1 and Kir6.2, confirmed by the reduced secretory response to tolbutamide (Figure 3C). These subunits of the ATP-sensitive K⁺ channel respond to changes in the intracellular ATP/ADP concentration ratio and thereby couple cellular metabolism to membrane electrical activity and insulin secretion. Studies of knockout mice have shown that the loss of ATP-sensitive K⁺ channels results in impairment of glucose-stimulated insulin secretion (33, 34). In addition, *Sur1* knockout mice exhibited increased basal plasma insulin levels, because the rate of return to the basal rate of insulin

secretion after a fall in blood glucose concentration was reduced, resulting in hypoglycemia during fasting that was similar to that observed in β PKC $\lambda^{-/-}$ mice. The ATP-sensitive K⁺ channel and hexokinases 1 and 2 have been shown to be targets of HNF3 β regulation in β cells (32, 35). Together, these various observations suggest that the downregulation of HNF3 β induced by PKC λ ablation is responsible for the changes in the expression of the genes for hexokinases 1 and 2, Sur1, and Kir6.2, and that these changes then result in the altered insulin secretion apparent in the islets of β PKC $\lambda^{-/-}$ mice. Expression of the *Glut2* gene was also reduced in islets of β PKC $\lambda^{-/-}$ mice. The Glut2 transporter mediates glucose uptake by pancreatic β cells, the first step in the signaling cascade responsible for glucose-stimulated insulin secretion. Ablation of Glut2 from mouse β cells resulted in loss of the initial phase of glucose-induced insulin release; the second phase was preserved but diminished (36), consistent with the phenotype of β PKC $\lambda^{-/-}$ mice.

To verify the role of PKC λ in insulin secretion suggested by our results, we examined the effects of restoration of expression of this enzyme with the use of an adenoviral vector in islets isolated from β PKC $\lambda^{-/-}$ mice. Restoration of PKC λ expression indeed normalized glucose-stimulated insulin secretion as well as the abundance of mRNAs for HNF3 β , hexokinase 1, and Kir6.2 in the β PKC $\lambda^{-/-}$ islets, consistent with our conclusion that PKC λ regulates the expression of genes whose products mediate the induction of insulin release by glucose. We also examined the effects of normalization of HNF3 β expression in islets derived from β PKC $\lambda^{-/-}$ mice. HNF3 β (Foxa-2) is a member of the hepatocyte nuclear factor 3 and forkhead family of transcription factors that also includes HNF3 α (Foxa-1) and HNF3 γ (Foxa-3). HNF3 proteins play important roles in maintaining glucose homeostasis through regulation of gene expression in the liver. HNF3 β is thought to act upstream of Pdx1, HNF1 α , and HNF4 α during pancreatic development. In mature β cells, HNF3 β regulates the expression of genes important for β cell function, including those for Pdx1, Nkx6.2, Sur1, and Kir6.2 (35, 37). A dominant negative form of HNF3 β was also shown to reduce the expression of the genes for Sur1 and Kir6.2 by 70% in INS-1 cells (32). The cells expressing this HNF3 β mutant also exhibited upregulation of *hexokinase 1* and *hexokinase 2* as well as failure to regulate *Pdx1*, consistent with our present results. In contrast to our results, however, expression of the HNF3 β mutant resulted in downregulation of *HNF1 α* and *HNF4 α* ; it is possible that the abundance of *HNF1 α* and *HNF4 α* mRNAs was unaffected in the islets of β PKC $\lambda^{-/-}$ mice because the amount of *HNF3 β* mRNA was reduced by only 40–50% and thus might still be sufficient to allow normal transcription of the *HNF1 α* and *HNF4 α* genes. Although mutations of the *HNF3 β* gene are not a common cause of maturity-onset diabetes of the young, a loss-of-function mutation (A86T) of this gene has been associated with type 2 diabetes (38), suggesting that HNF3 β gene mutations might confer predisposition to this disease. Indeed, normalization of HNF3 β expression in islets isolated from β PKC $\lambda^{-/-}$ mice resulted in a significant reversal of the impairment of glucose-induced insulin secretion. Taken together, we conclude that PKC λ plays a prominent role in the regulation of glucose-induced insulin secretion by modulating the gene expression of *HNF3 β* , *Glut2*, *hexokinase*, *Sur1*, and *Kir6.2*, and most likely also other effector molecules important for β cell function.

Ablation of the IGF-1 receptor in mouse β cells has been shown to impair insulin secretion, possibly as a result of a reduction in the levels of expression of *Glut2*, *glucokinase*, and *HNF3 β* (7). The β cell-specific IGF-1 receptor knockout mouse and our



β PKC $\lambda^{-/-}$ mice thus both show similar changes in insulin secretion and in gene expression. Silencing of the IGF-1 receptor by RNA interference also inhibited PI3K activity and glucose-stimulated insulin secretion in MIN6 cells and increased the extent of basal insulin release (9). Given that PI3K and 3-phosphoinositide-dependent kinase-1 contribute to the signaling pathway that links receptor tyrosine kinases to PKC λ (24, 29, 39), PKC λ might mediate signaling from the IGF-1 receptor to gene transcription in pancreatic β cells.

Hypersecretion of insulin at basal blood glucose levels and impairment of glucose-stimulated insulin secretion are characteristics of type 2 diabetes (40), and lipotoxicity manifests a similar phenotype of insulin secretion (41). Further elucidation of the intracellular signaling mediated by PKC λ in pancreatic β cells might thus provide a basis for the development of new therapeutic strategies for diabetes.

Methods

Animals and genotyping. Mice (PKC $\lambda^{lox/lox}$) harboring a “floxed” PKC λ gene were generated by homologous recombination (K. Akimoto et al., unpublished observations). Mice expressing the Cre recombinase under the control of the rat insulin 2 gene promoter were generated as described previously (18). The genetic background of mice was derived from a hybrid of C57BL/6, 129sv, and DBA-2 strains. Animals were maintained under a 12-hour-light, 12-hour-dark cycle and fed either normal chow or a high-fat diet from the time of weaning (3 weeks old) for a 21-week period. The high-fat diet contained 30% (wt/wt) fat (14% bovine fat, 14% porcine fat, 2% soybean oil) (42). Mice were genotyped by PCR analysis with primers (5' and 3', respectively) specific for the Cre transgene (5'-ATGTCCAATTTACTGACCG-3', 5'-CGCCGCATAACCAGTGAAC-3') or for the floxed PKC λ allele (5'-CATGCAGTGTACTGGCATAGCCACC-3', 5'-AGAGGCAGC-CAAAGCCCTGCTCTCC-3'). PCR was performed for 39 cycles of 94°C for 30 seconds, 50°C for 30 seconds, and 72°C for 45 seconds or for 35 cycles of 94°C for 30 seconds, 66°C for 30 seconds, and 72°C for 30 seconds, respectively. This study was performed according to the guidelines of the Animal Ethics Committee of Kobe University Graduate School of Medicine.

Immunoprecipitation and immunoblot analysis. For determination of the abundance of PKC λ and PKC ζ , total tissue homogenates were subjected to immunoprecipitation with rabbit polyclonal antibodies against PKC λ (α L190) or against PKC ζ (α Z170) (14), and the resulting precipitates were subjected to immunoblot analysis with mouse mAbs against PKC λ /t (Transduction Laboratories) or rabbit polyclonal antibodies against PKC ζ (Upstate Biotechnology Inc.), respectively. Goat polyclonal antibodies against HNF3 β obtained from Santa Cruz Biotechnology Inc. were used to determine the abundance of HNF3 β by immunoblot analysis.

Analysis of metabolic parameters. Blood samples were collected from the tail vein and plasma samples were separated by centrifugation of blood in a microcentrifuge for 5 minutes at 4°C. Blood glucose level was determined with a glucometer (Glutest Pro; Sanwa Kagaku Kenkyusho Co.). Plasma insulin concentration was measured with an ELISA kit with a mouse insulin standard (Shibayagi Co.) or by RIA with a rat insulin standard (Eiken Chemical Co.). All assays were performed in duplicate (43).

Intraperitoneal glucose tolerance and insulin tolerance test. Mice were deprived of food for 16 hours and then anesthetized with pentobarbital (30 mg per kilogram of body weight). Blood was collected immediately before as well as 30, 60, and 120 minutes after the intraperitoneal injection of glucose (1.5 mg/kg) (44). For insulin tolerance testing, mice in the randomly fed state were injected with 0.75 U/kg body weight of human regular insulin. Blood glucose and plasma insulin levels were measured as described above.

Assay of insulin secretion from isolated islets. Islets were isolated from 6-month-old mice by collagenase digestion and subsequent centrifugation

over a Histopaque (Sigma-Aldrich) gradient as described previously (44). For assay of insulin release, 5 islets were manually selected, incubated in Krebs-Ringer solution, and stimulated at 37°C with various concentrations of either glucose for 1 hour, KCl for 30 minutes, or tolbutamide for 1 hour. The islets were then collected by centrifugation, and the supernatant was assayed for insulin content by RIA as described above. For measurement of islet insulin content, islets were solubilized in acid-ethanol solution (74% ethanol, 1.4% HCl) overnight at 4°C before insulin RIA.

Immunostaining and morphometric analysis of islets. The pancreas was removed from 6-month-old mice, weighed, fixed overnight in Bouin's solution, and embedded in paraffin. Consecutive 4- μ m-thick sections were cut from the tissue and mounted on glass slides. After rehydration, the sections were stained with H&E. Immunostaining for insulin and glucagon was performed with guinea pig antibodies against insulin and rabbit antibodies against glucagon (DAKO Japan); immune complexes were detected with secondary antibodies conjugated with Cy3 or FITC, respectively (Jackson ImmunoResearch Laboratories Inc.). Immunostaining for PKC λ was performed with antibodies against PKC λ (α L190). For quantitation of β cell area, 4 animals of each genotype were analyzed at 6 months of age. Sections of paraffin-embedded pancreas were immunostained at 200- μ m intervals to avoid measurement of the same islets twice. Images of β cells and of the entire pancreas were obtained with a digital camera (EOS D30; Canon Inc.) and analyzed with the use of NIH Image 1.60 software, as described previously (45). The β cell area was expressed as a percentage of the total pancreatic area surveyed.

Electron microscopy. Two PKC $\lambda^{lox/lox}$ and 2 β PKC $\lambda^{-/-}$ mice were anesthetized with ether and subjected to intracardial perfusion with 2% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium phosphate buffer (pH 7.4). The pancreas was excised from each mouse, cut into small pieces, and immersed overnight in the same fixative. The tissue was then exposed to 2% OsO $_4$, treated with 2% uranyl acetate, dehydrated with ethanol, and embedded in Epon 812 (Nacalai Tesque). For light microscopy, 1- μ m-thick sections were cut and stained with toluidine blue. For electron microscopy, thin sections were stained with uranyl acetate and lead citrate before examination with a Hitachi 7100 electron microscope (Hitachi Ltd.).

Quantitation of mRNA by real-time RT-PCR. Total cellular RNA was isolated from islets of PKC $\lambda^{lox/lox}$ and β PKC $\lambda^{-/-}$ mice with the use of an RNeasy kit (QIAGEN Sciences). Real-time RT-PCR analysis of the total RNA pooled from 6 animals of each genotype was performed with an ABI PRISM 7900 Sequence Detection System (Applied Biosystems); 36B4 mRNA was used as an internal standard as previously described (27). Each reaction was performed in triplicate.

Infection of isolated islets with adenoviral vectors. Adenoviral vectors encoding β -galactosidase or wild-type mouse PKC λ have been described previously (46). Complementary DNA encoding human HNF3 β was kindly provided by M. Stoffel (Rockefeller University, New York, New York, USA), and an adenoviral vector encoding HNF3 β was generated with an adenovirus expression kit (Takara Shuzo Co.) as described previously (47). Isolated islets were maintained in RPMI 1640 medium that contained 11.1 mM glucose and was supplemented with 10% FBS. About 200 islets were infected with the adenoviral vectors at an MOI of 10 PFUs per cell. Experiments were performed 24 hours after infection.

Statistical analysis. Data are expressed as means \pm SE and were analyzed with ANOVA. A *P* value of less than 0.05 was considered statistically significant.

Acknowledgments

We thank M.A. Magnuson for Ins-Cre mice, T. Kitamura and T. Aizawa for helpful discussions, and K. Satomura and M. Kawasaki for technical assistance. This work was supported by grants from the Intellectual Cluster Formation Project, and from the 21st Century COE Program “Center of Excellence for Signal Transduction Disease:



Diabetes Mellitus as Model,” from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (to M. Kasuga).

Received for publication May 24, 2004, and accepted in revised form October 26, 2004.

Address correspondence to: Yoshiaki Kido, Department of Clinical Molecular Medicine, Division of Diabetes and Digestive and Kidney Diseases, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017, Japan. Phone: 81-78-382-5861; Fax: 81-78-382-2080; E-mail: kido@med.kobe-u.ac.jp.

- Rhodes, C.J. 2000. IGF-I and GH post-receptor signaling mechanisms for pancreatic beta-cell replication. *J. Mol. Endocrinol.* **24**:303–311.
- Withers, D.J., et al. 1999. Irs-2 coordinates Igf-1 receptor-mediated beta-cell development and peripheral insulin signalling. *Nat. Genet.* **23**:32–40.
- Leibiger, I.B., Leibiger, B., Moede, T., and Berggren, P.O. 1998. Exocytosis of insulin promotes insulin gene transcription via the insulin receptor/PI-3 kinase/p70 S6 kinase and CaM kinase pathways. *Mol. Cell.* **1**:933–938.
- Leibiger, B., et al. 2001. Selective insulin signaling through A and B insulin receptors regulates transcription of insulin and glucokinase genes in pancreatic beta cells. *Mol. Cell.* **7**:559–570.
- Persaud, S.J., Harris, T.E., Burns, C.J., and Jones, P.M. 1999. Tyrosine kinases play a permissive role in glucose-induced insulin secretion from adult rat islets. *J. Mol. Endocrinol.* **22**:19–28.
- Kulkarni, R.N., et al. 1999. Tissue-specific knockout of the insulin receptor in pancreatic beta cells creates an insulin secretory defect similar to that in type 2 diabetes. *Cell.* **96**:329–339.
- Kulkarni, R.N., et al. 2002. Beta-cell-specific deletion of the Igf1 receptor leads to hyperinsulinemia and glucose intolerance but does not alter beta-cell mass. *Nat. Genet.* **31**:111–115.
- Xuan, S., et al. 2002. Defective insulin secretion in pancreatic beta cells lacking type 1 IGF receptor. *J. Clin. Invest.* **110**:1011–1019. doi:10.1172/JCI200215276.
- da Silva Xavier, G., Qian, Q., Cullen, P.J., and Rutter, G.A. 2004. Distinct roles for insulin and insulin-like growth factor-1 receptors in pancreatic beta-cell glucose sensing revealed by RNA silencing. *Biochem. J.* **377**:149–158.
- Williams, M.R., et al. 2000. The role of 3-phosphoinositide-dependent protein kinase 1 in activating AGC kinases defined in embryonic stem cells. *Curr. Biol.* **10**:439–448.
- Tuttle, R.L., et al. 2001. Regulation of pancreatic beta-cell growth and survival by the serine/threonine protein kinase Akt1/PKB α . *Nat. Med.* **7**:1133–1137.
- Bernal-Mizrachi, E., Wen, W., Stahlhut, S., Welling, C.M., and Permutt, M.A. 2001. Islet beta cell expression of constitutively active Akt1/PKB α induces striking hypertrophy, hyperplasia, and hyperinsulinemia. *J. Clin. Invest.* **108**:1631–1638. doi:10.1172/JCI200113785.
- Pende, M., et al. 2000. Hypoinsulinaemia, glucose intolerance and diminished beta-cell size in S6K1-deficient mice. *Nature.* **408**:994–997.
- Selbie, L.A., Schmitz-Peiffer, C., Sheng, Y., and Biden, T.J. 1993. Molecular cloning and characterization of PKC iota, an atypical isoform of protein kinase C derived from insulin-secreting cells. *J. Biol. Chem.* **268**:24296–24302.
- Furukawa, N., et al. 1999. Possible involvement of atypical protein kinase C (PKC) in glucose-sensitive expression of the human insulin gene: DNA-binding activity and transcriptional activity of pancreatic and calphostin C-sensitive but phorbol 12-myristate 13-acetate (PMA) and Go 6976-insensitive pathway. *Endocr. J.* **46**:43–58.
- Harris, T.E., Persaud, S.J., and Jones, P.M. 1996. Atypical isoforms of PKC and insulin secretion from pancreatic beta-cells: evidence using Go 6976 and Ro 31-8220 as PKC inhibitors. *Biochem. Biophys. Res. Commun.* **227**:672–676.
- Buteau, J., et al. 2001. Protein kinase C ζ activation mediates glucagon-like peptide-1-induced pancreatic beta-cell proliferation. *Diabetes.* **50**:2237–2243.
- Postic, C., et al. 1999. Dual roles for glucokinase in glucose homeostasis as determined by liver and pancreatic beta cell-specific gene knock-outs using Cre recombinase. *J. Biol. Chem.* **274**:305–315.
- Cui, Y., et al. 2004. Essential role of STAT3 in body weight and glucose homeostasis. *Mol. Cell. Biol.* **24**:258–269.
- Tabuse, Y., et al. 1998. Atypical protein kinase C cooperates with PAR-3 to establish embryonic polarity in *Caenorhabditis elegans*. *Development.* **125**:3607–3614.
- Muller, H.A. 2000. Genetic control of epithelial cell polarity: lessons from *Drosophila*. *Dev. Dyn.* **218**:52–67.
- Doe, C.Q., and Bowerman, B. 2001. Asymmetric cell division: fly neuroblast meets worm zygote. *Curr. Opin. Cell Biol.* **13**:68–75.
- Suzuki, A., et al. 2001. Atypical protein kinase C is involved in the evolutionarily conserved par protein complex and plays a critical role in establishing epithelia-specific junctional structures. *J. Cell Biol.* **152**:1183–1196.
- Bjorkoy, G., Perander, M., Overvatn, A., and Johansen, T. 1997. Reversion of ras- and phosphatidylcholine-hydrolyzing phospholipase C-mediated transformation of NIH 3T3 cells by a dominant interfering mutant of protein kinase C lambda is accompanied by the loss of constitutive nuclear mitogen-activated protein kinase/extracellular signal-regulated kinase activity. *J. Biol. Chem.* **272**:11557–11565.
- Akimoto, K., et al. 1998. Atypical protein kinase C lambda binds and regulates p70 S6 kinase. *Biochem. J.* **335**:417–424.
- Suzuki, A., Akimoto, K., and Ohno, S. 2003. Protein kinase C λ /t (PKC λ /t): a PKC isotype essential for the development of multicellular organisms. *J. Biochem.* **133**:9–16.
- Matsumoto, M., et al. 2003. PKC λ in liver mediates insulin-induced SREBP-1c expression and determines both hepatic lipid content and overall insulin sensitivity. *J. Clin. Invest.* **112**:935–944. doi:10.1172/JCI200318816.
- Perander, M., Bjorkoy, G., and Johansen, T. 2001. Nuclear import and export signals enable rapid nucleocytoplasmic shuttling of the atypical protein kinase C λ . *J. Biol. Chem.* **276**:13015–13024.
- Akimoto, K., et al. 1996. EGF or PDGF receptors activate atypical PKC λ through phosphatidylinositol 3-kinase. *EMBO J.* **15**:788–798.
- Urban, R.J., Boedenberg, Y.H., Jiang, J., Denner, L., and Chedrese, J. 2004. Protein kinase C iota enhances the transcriptional activity of the porcine P450 side chain cleavage insulin-like response element. *Am. J. Physiol. Endocrinol. Metab.* **286**:E975–E979.
- Becker, T.C., BeltrandRio, H., Noel, R.J., Johnson, J.H., and Newgard, C.B. 1994. Overexpression of hexokinase I in isolated islets of Langerhans via recombinant adenovirus. Enhancement of glucose metabolism and insulin secretion at basal but not stimulatory glucose levels. *J. Biol. Chem.* **269**:21234–21238.
- Wang, H., Gauthier, B.R., Hagenfeldt-Johansson, K.A., Iezzi, M., and Wollheim, C.B. 2002. Foxa2 (HNF3 β) controls multiple genes implicated in metabolism-secretion coupling of glucose-induced insulin release. *J. Biol. Chem.* **277**:17564–17570.
- Seghers, V., Nakazaki, M., DeMayo, F., Aguilar-Bryan, L., and Bryan, J. 2000. Sur1 knockout mice. *J. Biol. Chem.* **275**:9270–9277.
- Miki, T., et al. 1998. Defective insulin secretion and enhanced insulin action in K $_{ATP}$ channel-deficient mice. *Proc. Natl. Acad. Sci. U. S. A.* **95**:10402–10406.
- Sund, N.J., et al. 2001. Tissue-specific deletion of Foxa2 in pancreatic beta cells results in hyperinsulinemic hypoglycemia. *Genes Dev.* **15**:1706–1715.
- Guillam, M.T., et al. 1997. Early diabetes and abnormal postnatal pancreatic islet development in mice lacking Glut-2. *Nat. Genet.* **17**:327–330.
- Lee, C.S., et al. 2002. Foxa2 controls Pdx1 gene expression in pancreatic beta-cells in vivo. *Diabetes.* **51**:2546–2551.
- Zhu, Q., et al. 2000. Identification of missense mutations in the hepatocyte nuclear factor-3 β gene in Japanese subjects with late-onset Type II diabetes mellitus. *Diabetologia.* **43**:1197–1200.
- Kotani, K., et al. 1998. Requirement of atypical protein kinase C λ for insulin stimulation of glucose uptake but not for Akt activation in 3T3-L1 adipocytes. *Mol. Cell. Biol.* **18**:6971–6982.
- Porte, D. 1991. Banting lecture 1990: beta-cells in type II diabetes mellitus. *Diabetes.* **40**:166–180.
- Unger, R.H. 1995. Lipotoxicity in the pathogenesis of obesity-dependent NIDDM. Genetic and clinical implications. *Diabetes.* **44**:863–870.
- Inoue, H., et al. 2004. Role of STAT-3 in regulation of hepatic gluconeogenic genes and carbohydrate metabolism in vivo. *Nat. Med.* **10**:168–174.
- Kido, Y., et al. 2000. Tissue-specific insulin resistance in mice with mutations in the insulin receptor, IRS-1, and IRS-2. *J. Clin. Invest.* **105**:199–205.
- Kitamura, T., et al. 2001. Preserved pancreatic beta-cell development and function in mice lacking the insulin receptor-related receptor. *Mol. Cell. Biol.* **21**:5624–5630.
- Kido, Y., et al. 2002. Effects of mutations in the insulin-like growth factor signaling system on embryonic pancreas development and beta-cell compensation to insulin resistance. *J. Biol. Chem.* **277**:36740–36747.
- Sakaue, H., et al. 1997. Phosphoinositide 3-kinase is required for insulin-induced but not for growth hormone- or hyperosmolarity-induced glucose uptake in 3T3-L1 adipocytes. *Mol. Endocrinol.* **11**:1552–1562.
- Kitamura, T., et al. 1998. Requirement for activation of the serine-threonine kinase Akt (protein kinase B) in insulin stimulation of protein synthesis but not of glucose transport. *Mol. Cell. Biol.* **18**:3708–3717.